

Assessment of Climate Risks and Adaptation Options

for Carpathian Forest Ecosystems and their Services

CARPATHIAN CONVENTION



Copyright © 2025 Carpathian Convention

This publication may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgement of the source is made. The Secretariat of the Carpathian Convention would appreciate receiving a copy of any publication that uses this publication as a source.

No use of this publication may be made for resale or any other commercial purpose whatsoever without prior permission in writing from the Secretariat of the Carpathian Convention.

Disclaimers

The views expressed in this publication are those of the authors and do not necessarily reflect the views or policies of the Carpathian Convention, contributory organizations, donor or any governmental authority or institution with which authors or contributors are affiliated and neither do they imply any endorsement. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the Carpathian Convention or contributing organizations concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Mention of a commercial company or product in this document does not imply endorsement by the Carpathian Convention or the authors. The use of information from this document for publicity or advertising is not permitted. Trademark names and symbols are used in an editorial fashion with no intention of infringement of trademark or copyright laws.

The opinions, figures and estimates set forth in this booklet are the responsibility of the authors and should not necessarily be considered as reflecting the views or carrying the endorsement of the Carpathian Convention. While reasonable efforts have been made to ensure that the contents of this publication are factually correct and properly referenced, the Carpathian Convention does not accept responsibility for the accuracy or completeness of the contents and shall not be liable for any loss or damage that may be occasioned directly or indirectly through the use of, or reliance on, the contents of this publication. We regret any errors or omissions that may unwittingly have been made.

© Photos and illustrations as specified.

Suggested citation: Keeton, W.S., McCallum, S., Kuraś, K., Fischer, E., & Egerer, H. (2025). Assessment of climate risks and adaptation options for Carpathian forest ecosystems and their services. Secretariat of the Carpathian Convention. Vienna, Austria. 65 pp.

Front page photo: Miloslav Brandt

ASSESSMENT OF CLIMATE RISKS AND ADAPTATION OPTIONS FOR CARPATHIAN FOREST ECOSYSTEMS AND THEIR SERVICES

AUTHORS

William S. Keeton
University of Vermont

Sabine McCallum
UN Environment Programme

Klaudia Kuraś
UN Environment Programme

Elena Fischer
UN Environment Programme

Harald Egerer
UN Environment Programme

EDITOR

Elena Fischer
UN Environment Programme


DESIGN AND LAYOUT

Elena Fischer
UN Environment Programme

ACKNOWLEDGEMENTS

We authors are grateful to the Carpathian Convention National Focal Points, members of the Workings Groups Forest and Climate Change, Science for the Carpathians, Trust for Mutual Understanding (New York, NY), and the United States Fulbright Scholars Program for supporting development of this assessment.

Photo: Mircea Verghelet



CONTRIBUTORS

Highly valued contributions about climate risks, impacts and adaptation options in the Carpathians was provided through a survey by:

CZECH REPUBLIC

Miroslav Svoboda

Czech University of Life Sciences Prague

Eliška Rolfova

Ministry of the Environment of the Czech Republic

Radek Pokorný

Mendel University in Brno

HUNGARY

Borbála Galos

University of Sopron

Imelda Somodi

Centre for Ecological Research

POLAND

Bożydar Neroj

Bureau for Forest Management and Geodesy

Wojciech Grodzki

Forest Research Institute

Małgorzata Czyżewska

Directorate General of the State Forest of Poland

ROMANIA

Laurentiu Radu

Ministry of Environment, Waters and Forest of Romania

Liliana Virtopeanu

Ministry of Environment, Waters and Forest of Romania

Borz Stelian Alexandru

Transilvania University of Brasov, Department of Forest Engineering

Păcurar Victor Dan

Transilvania University of Brasov

Sorin Cheval

National Meteorological Administration of Romania

Popa Ionel

Forest Research and Management Institute, Romania

SLOVAKIA

Libor Ulrych

State Nature Conservancy of Slovak Republic

SERBIA

Ilija Dordevic

Institute of forestry, Department for spatial planning, GIS and forest policy, Assistant director for international cooperation

UKRAINE

Liubov Poliakova

Head of International Cooperation, Science and Public Relation Division, State Forest Resources Agency

Volodymyr Korzhov

Deputy Head of Ukrainian Scientific Institute of Mountain Forestry

ABBREVIATIONS

CARPATCLIM	Climate in the Carpathian Region
CC	Carpathian Convention
CLIMAFORCEELIFE	Climate-Smart Forest Management for Central and Eastern Europe
COP	Conference of the Parties
CO2	Carbon dioxide
EbA	Ecosystem based Adaptation
EC	European Commission
EEA	European Environment Agency
EFI	European Forest Institute
EU	European Union
ETC-ULS	European Topic Centre on Urban, Land, and Soil Systems
FAO	Food and Agriculture Organization of the United Nations
FLAM	Wildfire Climate Impacts and Adaptation Model
FoRISK	Pan-European Forest risk knowledge mechanism
GEDI	Global Ecosystem Dynamics Investigation
GEOS	Global Earth Observation System of Systems
IIASA	International Institute for Applied Systems Analysis
IPCC	The Intergovernmental Panel on Climate Change
NbS	Nature-based Solutions
NCS	Natural Climate Solutions
N/A	Not applicable
SAFERS	Forest Fire Emergencies in Resilient Societies
SUPERB	Systemic solutions for upscaling of urgent ecosystem restoration for forest-related biodiversity and ecosystem services
UN	United Nations
UNESCO	United Nations Educational, Scientific and Cultural Organization
UNFCCC	United Nations Framework Convention on Climate Change
WWF	World Wild Fund for Nature

TABLE OF CONTENTS

Key Messages	i
Executive Summary	v
Chapter 1: Introduction.....	1
1.1 Background	2
1.2 Previous Regional and Pan-European Assessments	4
Chapter 2: Assessment Methodology	6
2.1 Introduction	7
2.2 Mandate for Conducting the Assessment	7
2.3 Workshop Approach	8
2.4 Survey and Literature Review Approach.....	11
Chapter 3: Climate Change Risks and Impacts on Carpathian Forest Ecosystems and their Services	12
3.1 Introduction	13
3.2 Survey Results and Literature Review on Key Risks and Impacts	15
3.3 Areas of Continuing Debate	23
Chapter 4: Adaptation Approaches	24
4.1 Introduction	25
4.2 Response Options Along Identified Key Topics	25
Chapter 5: Relevant Initiatives and Actors	40
5.1 Introduction	41
5.2 Relevant Projects	41
5.3 Relevant Tools.....	42
5.4 Relevant Research Initiatives	44
5.5 Relevant Policies.....	45
Chapter 6: Conclusions and Recommendations	46
6.1 Introduction	47
6.2 Opportunities and Pathways.....	48
6.3 Knowledge Gaps and Research Needs	50
6.4 Conclusion	51
References	52
Photo Credits.....	55
ANNEX 1: Nominated Experts Supporting the Assessment	56
ANNEX 2: Survey.....	57



KEY MESSAGES

CONTEXT

Forest ecosystems are a precious ecological, economic and cultural resource in the Carpathian region. The Carpathians are home to over 200 plant species that are endemic to only this region and contain Europe's largest concentration of virgin, quasi-virgin and natural forests. Because these forests are largely inaccessible and contain large, uninterrupted expanses, they provide refuge for populations of large European mammalian species, such as the lynx, river otter, grey wolf, woodland bison, red deer, moose and brown bear.

Carpathian forests are vulnerable to disrupted ecological processes, species interactions and overall ecosystem dynamics as a result of historical forest management and climate change. Carpathian forests bear the legacy of a long history of intensive production-driven management since the 19th century, making them more vulnerable to large-scale tree mortality and dieback. Extensive deforestation from large-scale timber extraction and clearing for agriculture was common, and even-aged, plantation style forest management practices homogenized the forest structure and composition.

Climate change superimposes additional stresses on species and ecosystems both directly and indirectly. Climate change directly effects plant physiology, phenology and reproductive success, and alters and often increases disturbance frequencies and severities. Rising temperatures, altered precipitation patterns and increased frequency of extreme weather events already present significant challenges to forests' health and functioning. Interactions among these stressors are likely to shift the structure and function of forest ecosystems, exacerbating risks and vulnerabilities and altering the mix of services and habitats forests provide.

IDENTIFIED CLIMATE RISKS

This assessment seeks to address these challenges by presenting several key climate change-related risks and impacts of greatest concern to experts in the Carpathian region. These concerns were identified through a Carpathian-wide survey of focal experts supported by a literature review. These risks and impacts pose significant threats to forest ecosystems and their crucial ecosystem services, including:

ALTERED DISTURBANCE REGIMES

Increased disturbance risks will accelerate overall rates of forest change, exacerbating other threats such as the spread of invasive species, shifts in species range, and loss of important habitats for biodiversity. More frequent disturbances create feedback loops that diminish the provisioning of critical ecosystem services, such as timber and non-timber resource production, carbon storage and hydrologic regulation.

DROUGHT AND WILDFIRES

Drought events and associated disturbance risks in Carpathian forests have become increasingly common and severe in recent years. Drought negatively impacts Carpathian forests' productivity, timber resources, susceptibility to disturbances, biodiversity and habitats, and climate regulating functions. Drought conditions can elevate the risk of forest fires in the Carpathians, not only causing immediate damage and loss of vegetation, but also contributing to long-term ecological disturbances and changes in forest structure.



CHANGES IN HYDROLOGIC REGIMES

Changes in hydrologic regimes are perceived as a major vulnerability within the region, as they interact with drought, disturbance risks and human impairment of watershed functioning.

While shifts in precipitation patterns vary across the region, some areas may experience altered timing, intensity and rainfall distribution. This can result in more frequent and intense rainfall events or prolonged droughts; the latter leading to decreased soil moisture levels, thus impacting the water supply for vegetation and forest productivity. Conversely, intense rainfall events can result in rapid runoff, severe soil erosion and possible chemical loading. There is a risk of increased flood frequency, although predictions carry high uncertainty and vary spatially throughout the region.

Large-scale disturbances, such as fire, bark beetle outbreaks and defoliating insects, will reduce water uptake by trees and reduce infiltration into soils, thereby increasing peak flows in rivers associated with high precipitation events.

Poorly designed forest roads, development and impervious surfaces are additional risk factors by changing the form of the hydrograph, meaning that water after precipitation events is directly delivered more rapidly and at greater volumes into streams, rivers and other surface waters. This increases peak flows and flood vulnerability after high precipitation events.

ADAPTATION RESPONSES

Various adaptation response options can be employed to enhance the resilience and sustainability of forest ecosystems in the Carpathians by addressing and coping with interlinked impacts and risks. With reference to the Carpathian Convention Protocol on Sustainable Forest Management, recommended key pathways for implementing climate-resilient forest management practices include:

PROTECTING AND CONSERVING VIRGIN AND NATURAL FORESTS by establishing and effectively managing protected areas, national parks and nature reserves to preserve intact forest ecosystems, maintain biodiversity and provide refuges for species. Such forests should be protected to also enhance the carbon sequestration and storage capacity of protected areas (*cf. Article 10*).

ENHANCING FOREST LANDSCAPE CONNECTIVITY through ecological corridors, which connect fragmented forest patches, enabling species movement and facilitating gene flow. Recommended practices include creating ecological corridors and conserving and restoring riparian zones, which can enhance connectivity and provide climate refugia for species (*cf. Article 16*).

MANAGING, PREVENTING AND RESTORING FOREST FIRES by developing national and regional early warning systems, improving fire suppression capabilities and promoting community-based fire management approaches. Towards such management, prescribed burning should be more broadly used to restore stand structures capable of supporting low intensity fire where this was historically operative in maintaining forest health, and adequate resources and training for forest fire management should be ensured to mitigate the impacts of wildfires on forests and communities (*cf. Article 14 lit.3*).

RESTORING FORESTS AND REFORESTING by restoring and replanting degraded forest areas; converting non-endemic monocultures to site-specific endemic species compositions; broadly using close-to-nature silvicultural practices; promoting natural forest regeneration; and protecting and reintroducing rare native tree species in their natural ecosystem. Such response options for adaptive management can be implemented individually or in combination with others (*cf. Article 13; Article 8*).

PROMOTING SUSTAINABLE WOOD UTILIZATION AND DEVELOPING VALUE CHAINS FOR FOREST PRODUCTS, including supporting responsible harvesting practices, local processing industries and the use of sustainably sourced wood products (*cf. Article 9*).

ADOPTING ECOSYSTEM-BASED ADAPTATION that recognizes and harnesses the benefits provided by intact ecosystems, and promote sustainable livelihoods in an inclusive approach (*cf. Article 12; Article 14; Article 15*).

INCORPORATING TRADITIONAL AND LOCAL KNOWLEDGE in adaptation planning to gain valuable insights into forest dynamics, species behavior and ecosystem responses to climate change. Community involvement also promotes local ownership, improves livelihoods and fosters sustainable forest practices (*cf. Article 15*).

IMPLEMENTING EDUCATION AND CAPACITY BUILDING EFFORTS, such as providing training, education, and awareness campaigns for forest managers, policymakers, local communities, and relevant stakeholders to enhance understanding of climate change impacts, adaptation strategies and the importance of sustainable forest management (*cf. Article 15*).

Carpathian Convention Protocol on Sustainable Forest Management Referenced Articles:

- Article 8** *Maintaining and enlarging forest cover*
- Article 9** *Ensuring the productive functions of the forest and their role in rural development*
- Article 10** *Identification and protection of natural, especially virgin forests*
- Article 12** *Improvement of the protective forest functions*
- Article 13** *Promotion of restoration of close-to-nature forests*
- Article 14** *Forestry and climate change*
- Article 15** *Social function of forests*
- Article 16** *Forestry, wildlife and ecological networks*
- Article 17** *Compatible monitoring and information systems*
- Article 18** *Coordinated scientific research and exchange of information*



Fostering collaboration and transboundary cooperation among the Carpathian countries will be vital for effective climate change adaptation in forestry, especially through sharing knowledge and experiences with various approaches as well as developing and implementing pathways for addressing common challenges.

STRENGTHENING RESEARCH AND CLOSING KNOWLEDGE GAPS around climate-resilient pathways will enhance evidence-based decision-making for adaptation in the Carpathian forestry sector. Current knowledge gaps exist in the following topics:

- Developing capacity for improved regional-scale forest monitoring (*cf. Article 17*). Harmonizing monitoring programs and sharing data across borders would facilitate both coordinated adaptation, such as multilateral plans for assisted species migration, and enable comparison of research results across the region. Forest monitoring could also include an additional layer for forest ecosystem dynamics under climate change, including changes in forest structure, species distribution patterns, and ecosystem functioning. Furthermore, forest ecosystems can exhibit ecological thresholds, beyond which they may experience rapid and irreversible changes in structure and function.
- The importance of genetic diversity in forest ecosystems for adaptation, i.e. studying the genetic characteristics of tree species, assessing the adaptive potential of different genetic lineages, and investigating how genetic diversity influences ecosystem resilience and productivity.
- The socio-economic dimensions of adaptation in forestry. This includes assessing the economic viability and costs associated with different adaptation approaches, understanding the social acceptability and equity implications of adaptation interventions, and considering the impacts of adaptation on local communities and livelihoods.
- Assessing the effectiveness of various adaptive silviculture practices in Carpathian forests, i.e. evaluating the effects of different adaptation approaches on forest resilience, productivity, biodiversity and socio-economic factors. Long-term monitoring of adaptive practices will also be important to continuously (re-)evaluate their success. Research is further needed to investigate the interactions and synergies among different adaptation options. Understanding how different approaches can complement or conflict with each other is crucial for optimizing adaptation strategies and avoiding unintended consequences (*cf. Article 18*).

EXECUTIVE SUMMARY

The Carpathian Mountain region touches seven European countries and encompasses natural ecosystems of global ecological and cultural value. Carpathian forests harbour high levels of endemic biodiversity; contribute to local and national economies through both timber and non-timber products; generate life-sustaining ecosystem services, including climate regulation and flood control; and are central to the rural character and cultural heritage that have defined the region for centuries.

Yet these resources and services are vulnerable to the effects of climate change, exacerbated by forest mismanagement practices that have simplified forest structure and composition across vast areas.

The resulting landscape conditions interact with climatic trends of temperature increase, particularly during summer and in western portions of the region, and increasingly variable seasonal precipitation. The latter has become more variable year to year, exhibiting an overall increase in winter and autumn rainfall and a decline in summer precipitation.

Perceptions of increasing risk have been confirmed by several recent pan-European assessments. These have identified specific vulnerabilities facing forest resources in Carpathian countries, including climate induced physiological and reproductive stresses in trees, making them susceptible to diseases, the spread of invasive species, extreme climate events like heavy rainfall and floods, heatwaves, drought and wildfires, and altered natural disturbance regimes.



In response, the Carpathian Convention Conference of the Parties at its 6th meeting (COP6), held in 2020, called for an assessment of the impacts of climate change on the Carpathian forests and their ecosystem services

by relevant Carpathian Convention Working Groups and partners and with support of the Convention Secretariat. Through a series of workshops, conference sessions, online discussions, and a survey sent to focal experts nominated by the National Focal Points, the Secretariat obtained several key climate change-related risks and impacts of greatest concern to experts in the Carpathian region. This data forms the basis for the information presented in this assessment.

The survey was organized around seven themes identified by a subgroup of the Carpathian Convention Working Group on Sustainable Forest Management and the Working Group on Climate Change with stakeholder input as important topics to address. Focal experts (survey respondents) provided detailed information on key risks, climate impacts, and adaptation options of greatest concern within environmental and forest sectors based on these seven themes:

- 1. Forest growth & productivity**
- 2. Biomass & carbon stocks**
- 3. Tree mortality**
- 4. Changes in species range & abundance and habitat shifts**
- 5. Invasion by non-native species**
- 6. Forest ecosystem services**
- 7. Forest-water interactions, including hydrologic regulation & riparian dynamics**



To evaluate the qualitative data from survey responses and rank the specific risks by the number of mentions, a triangulation method was used for each theme. A similar process was used to extract and collate the priority adaptation options described by the focal experts in the survey. This process resulted in the synthesis of key risks and adaptation approaches presented in this assessment. Synthesized survey responses were supplemented with information from peer-reviewed scientific literature and interviews with key research groups at institutions located both within Carpathian countries and elsewhere in Europe. The literature and interviews echo the identified risks and impacts articulated by focal experts, as these concerns are also the subject of ongoing investigations at research institutions.

Based on the seven priority topics, the following six priority risks of concern were identified by focal experts:

- 1. Altered disturbance regimes**
- 2. Altered hydrologic regimes & flood risks**
- 3. Invasive species**
- 4. Changes in forest growth & productivity**
- 5. Altered species composition & distribution**
- 6. Feedback mechanisms & effects on ecosystem services, including carbon storage**

The most frequently mentioned risk to all seven themes was the increasing frequency and severity of natural disturbances, particularly insect outbreaks, fungal pathogens, and windstorms. Alteration of disturbance regimes caused by the interaction of climate change and human-modified forest landscape conditions (i.e. that increase vulnerability) will have cascading effects on ecosystem services, including timber and non-timber resource production, carbon storage, and hydrologic regulation.

Drought and related forest fire risks were the second-most mentioned, illustrating their great regional importance given recent climate trends and severe drought events. Drought was frequently cited by focal experts as posing grave consequences for forest growth and productivity, regional tree mortality rates, biodiversity, future shifts in species composition and the climate regulating functions of Carpathian forests.

Additional risks and impacts identified by focal experts varied. These included flood risks and their connection to forest cover, forest roads, and forest management; spread of invasive insect pests, tree pathogens, and noxious plants; range shifts in the distribution of biodiversity; and concerns over increased land-use pressures on forest ecosystems.

Focal experts had different views on some issues, such as restoration of older forests, the carbon sequestration and storage value of older forests, and whether forest management intensity should be increased or decreased to mitigate disturbance risks.

The assessment identifies several key pathways for climate-resilient forest management practices in the Carpathians. Adaptation approaches for enhancing the resilience and sustainability of forest ecosystems in the face of climate impacts and risks identified by focal experts included the following:

- **Adaptive silviculture**
- **Afforestation & reforestation**
- **Carbon forestry**
- **Rewilding & core area protection**
- **Increase resilience to disturbance**
- **Assisted dispersal & migration**
- **Management practices to maintain/improve the ability of forests to resist pests and pathogens**
- **Close-to-nature silviculture & ecosystem-based management**
- **Integrating forest-water interactions into adaptive forest management**

The most frequently mentioned adaptation response was forest restoration, while sustainable management—including broader use of close-to-nature silviculture and continuous cover forestry—was the second-most mentioned response.

Adaptation responses like forest restoration, afforestation and reforestation, and silviculture practices reduce drought stress, while conversion cutting to restore endemic species composition creates more heterogeneous landscapes with enhanced resilience to disturbance and drought.

Focal experts frequently mentioned restoration as a key adaptation pathway, specifically the need for climate-adapted regeneration practices.


There is great consensus between the focal experts and the scientific literature regarding the adaptive value in close-to-nature silviculture, continuous cover forestry, riparian buffers and upgraded forest road

engineering, expanding protected areas networks, and protecting remaining virgin forests specifically.

The assessment describes ongoing initiatives that complement the above-mentioned climate adaptation pathways. These include pan-European initiatives to map, monitor, and better predict current and future forest disturbance risks; recently published databases on European primary forests; initiatives that individual Carpathian countries have

taken to protect natural and virgin forests; growing interest in and demonstration of “climate-smart” forestry practices; and wide-scale tree planting programs.

Together with regional harmonization of climate risk monitoring and forecasting, accelerated investment in climate adaptation pathways would offer great promise for sustaining Carpathian forest ecosystems and their vital services into the future.





1

Introduction



Figure 1. Visual representation of the Carpathian mountain region.

1.1 BACKGROUND

The Carpathian Mountains are the backbone of Central and Eastern Europe, forming an arc of about 1,500 km and covering an area of about 209,000 km². Stretching across seven countries, the mountain range extends from eastern Czech Republic, through Hungary, Poland, Romania, Serbia, Slovakia and Ukraine.

Partly because of their history as the frontier between shifting geopolitical boundaries, the Carpathians harbour biological and cultural resources found nowhere else in Europe, including a wealth of natural habitats and Europe's largest concentration of "natural forests" (FAO definition¹) outside of Fennoscandia (Sabatini et al. 2018).

For this reason, the Carpathian region is recognized as having global cultural and ecological significance by the United Nations Educational, Scientific and Cultural Organization (UNESCO). Large, uninterrupted and inaccessible expanses of forests provide refuge for populations of large European mammal species, such as the lynx, river otter, grey wolf, woodland bison, red deer, moose and brown bear. Found only in this region are over 200 species of endemic plants and stands of virgin² European beech forests with areas larger than 10,000 ha. Subranges like the Făgăraș in Romania contain the largest areas of virgin or primary (never cleared) montane forests remaining in temperate Europe.

Due to the geographical remoteness of some areas and strong cultural identities, traditional rural agrarian lifestyles lost to other parts of Europe have been preserved in this region. These traditional practices include communal forest management systems in parts of western Ukraine and Transylvania in Romania.

Forests provide myriad ecosystem services that contribute to societal well-being, for example, by regulating water flows, protecting soils and conserving biodiversity. More than half of the Carpathians are forested, with stable or slightly increasing forest cover due to sustainable forestry practices, active reforestation, and natural forest succession on abandoned (often marginal) agricultural lands. All Carpathian countries regulate forest regeneration following timber harvest, and some countries prohibit deforestation and discourage forest expansion onto areas previously used as pasturelands.

However, despite the stable forest cover, Carpathian forests are increasingly vulnerable to numerous disturbances and anthropogenic pressures, raising concerns over their long-term sustainability, productivity, and functioning.

¹ Forest is determined both by the presence of trees and the absence of other predominant land uses. The trees should be able to reach a minimum height of 5 meters *in situ*. Source: Global Forest Resources Assessment 2020. @ FRA 2023.

² Synonymous with "primary," see Sabatini et al. (2018)

HISTORICAL FOREST MANAGEMENT

Carpathian forests have been profoundly altered by centuries of intensive, production-driven management, making them vulnerable to large-scale tree mortality and dieback. Intensive production-driven forestry systems originated before the 19th century and continued during the communist period. Lowland temperate forests were extensively logged under the Austro-Hungarian regime, while much of the native beech (*Fagus sylvatica*) and mixed species forests were converted to Norway spruce (*Picea abies*) plantations managed using even-aged silvicultural systems (Keeton et al. 2013). Spruce is native to the Carpathians but genetic varieties from outside the region were widely planted on sites where it was not historically endemic. These practices homogenized the structure and composition of forest stands and simplified landscape-level diversity, patterns, and patch mosaics, creating the spruce-dominated expanses seen in many areas today (Stoyko 1998).

The effects of climate change are likely to interact with these pre-existing vulnerabilities. For example, the homogenization of forest stands makes them more vulnerable to large-scale tree mortality and dieback from insects, fungal diseases, mistletoe and drought, which sometimes interact with other disturbances, like windstorms and forest fires, increasing tree mortality and disturbance contagion or spread (Koral et al. 2022).

CLIMATE CHANGE MITIGATION & CHALLENGES

This sets the stage for climate change, which superimposes additional stresses through direct effects on plant physiology, phenology, and reproductive success as well as indirect effects through altered and often increased disturbance frequencies and severities.

While forests have great potential in mitigating climate change impacts, altered natural disturbance regimes may limit or reverse their capacity to do so. Carpathian forests, and forests worldwide, are increasingly valued as Natural Climate Solutions (NCS), meaning that forests are potential solutions for reducing greenhouse gas emissions and storing carbon. For example, some regions, such as western Ukraine, have been a net carbon sink in recent decades due to forest recovery following post-Soviet farmland abandonment (Kuemmerle et al. 2011). However, increases in the frequency and severity of bark beetle outbreaks, forest fires and wind events have reduced carbon storage capacity in European forests, limiting or reversing the potential for forests to be NCS (Seidl et al. 2014).

Carpathian forest ecosystem dynamics will continue to change as compounded stresses from climate disruption, invasive species, land use pressures, and other factors increase, according to several recent pan-European assessments (see Table 1). With changes in ecosystem dynamics come alterations in the mix of ecosystem goods and services those forests provide.

Because much of the Carpathian region is dependent on natural resources for sustaining rural livelihoods, it is crucial to understand interactions between socioeconomic changes and climate change to build pathways for sustainable development and resilient forest ecosystems. Demographic change and increasing development pressures in the Carpathian region create complex challenges for addressing socioeconomic resilience to climate change.

Despite the high demand for traditional cultural and natural assets from the rest of Europe, the Carpathian region is struggling economically, as evidenced by the high level of unemployment and emigration towards western European countries. This starkly contrasts with certain locales experiencing increasing development pressure to expand tourism like ski areas and small-scale hydroelectric infrastructure.

Therefore, sustainable economic development and adaptive forest management approaches that increase resilience to climate-induced risks must go hand in hand.

Foresters, scientists and policy makers are challenged to integrate knowledge from multiple disciplines in addressing questions of climate change. However, in order to realize adaptive forest management approaches, key risks associated with climate change must first be identified, in addition to differentiating between natural climate variability and human-caused climate change and understanding their interactions.

Towards these aims, the next section provides an overview of previous relevant climate assessments from which this current assessment builds upon for guiding adaptive forest management.





1.2 PREVIOUS REGIONAL AND PAN-EUROPEAN ASSESSMENTS

Several European-wide and Carpathian-specific assessments conducted over the last decade aim to help guide ongoing climate mitigation and adaptation efforts in the Carpathian region (see Table 1). These assessments provided the launching point for this assessment and contain a wealth of information on a wide range of vulnerabilities, risks and adaptation options in the forest sector.

These previous assessments mapped and identified vulnerabilities to climate change across a wide range of environmental infrastructures and natural resources. They focused heavily on disturbance risks to forest resources, biodiversity and ecosystem services, like carbon sequestration and flood control. They describe a region and continent where forests are already experiencing climate change impacts and where disturbance regimes, biodiversity and resource provisioning will continue to change dramatically into the future. All these findings coincide with the expert opinions elucidated in the survey responses.

Table 1. Previous assessments of climate change vulnerability, risks, and adaptation for forests in Europe generally and the Carpathian region specifically.

PUBLICATION DATE	ASSESSMENT TITLE	GEOGRAPHIC SCOPE	CITATION
2014	Climate change vulnerability and ecosystem-based adaptation measures in the Carpathian region	Carpathians	Saskia et al. (2014)
2014	Future imperfect: climate change and adaptation in the Carpathians	Carpathians	Werners et al. (2014)
2016	Climate change adaptation in the Carpathian Mountain Region	Carpathians	Werners et al. (2016)
2017	Outlook on climate change adaptation in the Carpathian Mountains	Carpathians	Alberton et al. (2017)
2018	National climate change vulnerability and risk assessments in Europe, 2018	Compilation of national-level data for 33 countries	Füssel et al. (2018)
2020	Adaptation to climate change in sustainable forest management in Europe	Pan-European	Lindner et al. (2020)
2021	Vulnerability of European forests to natural disturbances	Pan-European	Forzieri et al. (2021)
2021	European forests for biodiversity, climate change mitigation and adaptation	Pan-European	Science for Environment Policy (2021)

Changing climatic conditions have been evident over the past 50 years and are highly likely to continue over the coming decades in the Carpathian region (Werners et al. 2014). Mean annual temperatures have increased by 1° to 1.5°C, exhibiting distinct seasonal and sub-regional trends, such as greater warming in summer and western portions of the region (Hlásny et al. 2017). Seasonal precipitation has become more variable year to year, exhibiting an overall increase in winter and autumn rainfall and a decline in summer precipitation (Ceballos et al. 2018). Rainfall patterns have also changed spatially throughout the Carpathian region, increasing in most places while decreasing in western and southeastern sub-regions (Spinoni et al. 2014).


According to a synthesis by Werners et al. (2014), regional climate models project a warmer future with more erratic precipitation, likely leading to more frequent and more intense heat waves and, idiosyncratically, both droughts and floods in the Carpathians. Mean annual temperature will increase 3.0-4.5°C from pre-industrial levels by the year 2100 (see Werners et al. 2014, page 5). The predicted warming intensity varies depending on the climate scenario. These comprise different pathways of societal response and greenhouse gas (GHG) emissions reduction (IPCC 2018). Models predict that total annual precipitation over this time period will increase by 300–400 mm in

northern portions of the region, while southern portions will experience a 100–150 mm decrease (see Werners et al. 2016, page 1). However, there is likely to be pronounced seasonal variation, with a 20% decrease in summer rainfall and a wintertime increase of 5-20% regionally (Werners et al. 2016).

These observed and predicted climatic trends, together with the ongoing trend of increased frequency of extreme weather events causing droughts and floods, are the ultimate drivers of the specific risks and impacts to Carpathian forests.

However, a range of adaptive forest management responses are available to help make forest ecosystems more resilient to climate risks. Several of the previous assessments make critical recommendations for policy and governance, providing a roadmap for Carpathian countries to coordinate adaptation planning and monitoring both regionally and at the European level.





2

Assessment Methodology

2.1 INTRODUCTION

Despite previous assessments of climate vulnerabilities for both the Carpathian region specifically and Europe as a whole, expert input is needed for ongoing regional climate adaptation work under the Carpathian Convention and its Working Groups on climate change and sustainable forest management.

This qualitative assessment aims to answer, “What are the climate risks and adaptation options of greatest concern

and interest to experts in each of the Carpathian countries?” By presenting the views of regional experts, this assessment seeks to better inform adaptive forest management and conservation planning at local, national, and regional levels.

The process of this assessment was initiated by the Carpathian Convention and entailed eliciting expert input on key concerns through a workshop and survey. Survey results were

cross-checked against a review of previous European-scale and regional-scale scientific assessments, expert interviews, and a literature review. Analysis of key risks and impacts and potential forest management strategies was conducted by using a triangulation method to identify areas of commonality in survey responses (see Chapter 2.4 below).

2.2 MANDATE FOR CONDUCTING THE ASSESSMENT

The Framework Convention on the Protection and Sustainable Development of the Carpathians (Carpathian Convention) is a subregional treaty that unites the seven Carpathian countries to achieve the environmental protection and the sustainable development of the Carpathians. Considering climate change’s increasingly adverse impacts, the Carpathian Convention’s governing body, the Conference of the Parties, and several other Convention entities recognized the need for strengthening protection of Carpathian forests.

At the 6th meeting of the Conference of the Parties to the Carpathian Convention (COP6) in 2020, the Parties adopted decisions encouraging the development of an assessment of the impacts of climate change on the Carpathian forests and their ecosystem services.³ Such an assessment was to be conducted by relevant Convention Working Groups and partners and facilitated by the Convention Secretariat.

Subsequently, the development process for this assessment was integrated into other Convention mechanisms as a concrete activity for achieving set objectives, furthering the momentum. The Carpathian Convention Working Group on Climate Change included the development of this assessment within their “Implementation Framework 2030 accompanying the Long-term Vision towards combating climate change in the Carpathians” and their “Workplan for the implementation period 2021-2023,” which outlines tangible activities and expected results for achieving the strategic objectives and related targets of the Long-term Vision 2030 towards combating climate change in the Carpathians.

³ *DECISION COP6/13 paragraph 5: Appreciates the strengthened cooperation between the WG Forest and the WG Climate Change and WG Biodiversity...welcomes the idea of collecting information from the Parties with the goal of assessing the impacts of climate change on the Carpathian forests and their ecosystem services, including, if possible, climate change effects on large carnivores and their habitats...and requests the relevant Working Groups and partners to support the development of such assessment, and the Secretariat to facilitate the process.*

DECISION COP6/18 paragraph 8: Specifically encourages the WG Forest and the WG Biodiversity and partners to jointly further develop with the WG Climate Change an assessment of the impacts of climate change on the Carpathian forests and their ecosystem services, including, if possible, climate change effects on large carnivores and their habitats, and requests the Secretariat to facilitate the process.

NOV 2020

6TH MEETING OF THE CONFERENCE OF THE PARTIES TO THE CARPATHIAN CONVENTION

2.3 WORKSHOP APPROACH

At the 8th meeting of the Working Group on Climate Change held online on 6 May 2021, Working Group members decided that engagement for the assessment of climate change impacts on Carpathian forests should begin at the Forum Carpaticum 2021.

Forum Carpaticum is a science conference for the Carpathian region organized by the Science for the Carpathians, a network of scientists, practitioners and students for sharing research and discussing challenges and solutions to ongoing environmental, economic and socio-political crisis in the Carpathian region. The Science for the Carpathians collaborates closely with the Carpathian Convention to help inform scientific-based decision-making by defining research priorities and sharing findings and recommendations with the Convention.

During the Forum Carpaticum on 22 June 2021, a special session and workshop on “Forest ecosystem vulnerabilities to climate change in the Carpathians” was organized online by Dr. William Keeton, University of Vermont professor and member of the Science for the Carpathians and Carpathian Convention Secretariat.⁴

MAY 2021

8TH MEETING OF THE WORKING GROUP ON CLIMATE CHANGE

JUNE 2021

FORUM CARPATICUM SPECIAL SESSION AND WORKSHOP

Workshop participants identified a variety of climate change impacts and risks of concern relative to forest ecosystems and resources by responding to the question, “**What information would you most like to see with an assessment of climate change impacts on the Carpathian forests and their ecosystem services that would be most beneficial for your country/work?**”

Their responses are illustrated as a word cloud in Figure 2, generated by the Mentimeter application.

Figure 2. Mentimeter-generated word cloud of climate change impacts and risks of greatest concern to participants in a workshop conducted at the Forum Carpaticum 2021.

While not a scientific approach, the word cloud was useful in visualizing the diverse topics of greatest concern for the workshop participants, with each topic weighted by the number of times it was mentioned as a rough indication of priority.

⁴ All workshop presentations and the final Workshop Report are accessible on the Carpathian Convention website.

The subsequent exchange and discussion at the workshop highlighted several additional areas of concern, which were collected and condensed into a list of sub-topics, priority risks and proposed adaptation responses. The word cloud results were also organized into an abbreviated list of topics by collating sub-themes around broader themes, such as ecosystem services, biodiversity or disturbances.

Following the workshop, a dedicated informal expert subgroup of the Working Group on Climate Change and the Working Group on Sustainable Forest Management was established, made up of experts nominated by the Carpathian Convention National Focal Points (*Annex 1: Nominated experts supporting the assessment*).

The subgroup aimed to support the development of the assessment while strengthening cooperation between these topics under the Carpathian Convention, in part fulfilling the COP6 Decisions and the “Long-term Vision 2030 towards combating climate change in the Carpathians.”

On 16 November 2021, the first expert subgroup meeting was conducted online, where members provided valuable input and background information and discussed key topics, impacts and risks facing forest ecosystems and adaptation responses in the Carpathian region.

The results and condensed lists from the Forum workshop were presented and considered, resulting in the final list of seven priority topics and matrix presented in Table 2. The Secretariat also presented a draft structure for the assessment to be populated with in-depth information collected from focal experts through a survey (*Annex 2: Survey*).

NOV 2021
1ST MEETING OF THE EXPERT SUBGROUP

Table 2. Key topics, impacts, and adaptation strategies identified as priorities by members of the expert subgroup of the Working Group on Climate Change and the Working Group on Sustainable Forest Management

PRIORITY TOPIC	POTENTIAL IMPACTS (SUMMARY OF INPUTS)	POTENTIAL ADAPTATION RESPONSES (SUMMARY OF INPUTS)
FOREST GROWTH & PRODUCTIVITY	<p>Increased physiological stress leading to reduced growth and vigor, including:</p> <ul style="list-style-type: none"> • Effects of temperature increase, variations in precipitation • Short term and long term effects of CO2 fertilization effects; interactions with stressors • Effects of altered disturbance regimes • Effects of drought 	<ul style="list-style-type: none"> • Adaptive silviculture • Pros and cons of expanded use of exotic species • Stocking management, including thinning to reduce stand densities
BIOMASS & CARBON STOCKS	<p>Changing/reduced carbon uptake and carbon dynamics (sequestration, storage, and fluxes), including:</p> <ul style="list-style-type: none"> • Effects of increased fire frequency and other disturbances due to climate change • Sensitivity of ecosystem respiration to rising temperatures 	<ul style="list-style-type: none"> • Improved forest management/carbon forestry • Afforestation/reforestation • Avoided land-use conversion • Core area protection/rewilding • Managing land use, fire, and other disturbances and non-climatic stressors

Table continued on next page

PRIORITY TOPIC	POTENTIAL IMPACTS (SUMMARY OF INPUTS)	POTENTIAL ADAPTATION RESPONSES (SUMMARY OF INPUTS)
TREE MORTALITY	<p><i>Interaction between mortality and disturbance risks, including:</i></p> <ul style="list-style-type: none"> • Effects of drought on mortality • Feedback between dead trees and forest fires: direct climate effects on increased risk of extreme events and forest fires; indirect effects from insect outbreaks, drought, disease processes, etc. 	<ul style="list-style-type: none"> • Adaptive management of fire, pests, and pathogens (variable approaches and opinions) • Restoration of site-endemic species • Restoration of landscape heterogeneity
CHANGES IN SPECIES RANGE & ABUNDANCE AND HABITAT SHIFTS	<p><i>Interaction between climatic factors and anthropogenic pressures, including:</i></p> <ul style="list-style-type: none"> • Extinction risk for species with intrinsically low dispersal rate and species in isolated habitats such as mountain tops • Habitat shifts • Impacts on keystone and flagship species and the composition of forest communities 	<ul style="list-style-type: none"> • Reduction of habitat modification and fragmentation, pollution, over-exploitation, and invasive species • Protected area expansion, assisted dispersal and migration, ex situ conservation
INVASION BY NON-NATIVE SPECIES	<p><i>Vulnerability of ecosystems to non-native species invasion due to disruptions of species interactions and altered climatic conditions</i></p> <ul style="list-style-type: none"> • Effects of biome shifts (as an extreme result of the above) upon the spectrum of ecosystem services provided 	<ul style="list-style-type: none"> • Forest management practices that reduce susceptibility to invasive species, largely based on reducing other stresses (except from climate) including expanded invasive species monitoring and control efforts
FOREST ECOSYSTEM SERVICES	<p><i>Effects of altered critical ecosystem services, including:</i></p> <ul style="list-style-type: none"> • Supporting services and primary production (timber and non-timber resource production) • Habitat and services provisioning (bioenergy, water) • Regulating services (climate regulation, hydrologic regulation, flood control, pollination, pest and disease control, carbon sequestration and storage) • Cultural services, including recreation and aesthetic and spiritual benefits 	<ul style="list-style-type: none"> • Adaptive forest management to build resilience within at-risk ecosystems by identifying the full set of drivers of change and most important areas and resources for protection and restoration • Foster inclusion of climate change considerations into the management of protected areas (incl. Natura2000) and core area restoration • Broad stakeholder and community engagement to build consensus on adaptation approaches • Socio-economic inclusive approaches that may also have community and cultural benefits (Ecosystem-based Adaptation)
FOREST-WATER INTERACTIONS, INCLUDING HYDROLOGIC REGULATION AND RIPARIAN DYNAMICS	<p><i>Effects of altered hydrology regimes upon forests' provision of watershed services, including:</i></p> <ul style="list-style-type: none"> • Impacts on water quality, aquatic habitats and species and soil resources • Reduced water uptake by trees and reduced soil infiltration due to large-scale disturbances, such as fire, bark beetle outbreaks, mistletoe and defoliating insects, leading to increased runoff, severe erosion and chemical loading • Accelerated nutrient cycling rates due to warmer temperatures, promoting increased forest growth and elevated nitrogen levels in streams 	<ul style="list-style-type: none"> • Better integrate water-related ecosystem services supply into climate-smart forest management objectives • Broader adoption of riparian buffer standards • Improved forest road planning, design, and regulation

2.4 SURVEY AND LITERATURE REVIEW APPROACH

The Secretariat together with Dr. William S. Keeton developed and administered a survey (*Annex 2: Survey*) to 18 focal experts from universities, research institutes, governmental ministries and state forest agencies (*Annex 1: Nominated experts supporting the assessment*). The intent was to gather detailed qualitative data on key risks and adaptation responses according to the seven priority topics of concern developed by the expert subgroup (see Chapter 2.2). The survey contained a series of questions following the structure of Table 3, where focal experts filled in the matrices for each of the seven priority topics, along with answering other related survey questions. They were encouraged to consult with colleagues for gathering further input.

Survey responses were received by the Secretariat in the first half of 2022. Survey results were cross-checked against a review of previous European-scale and regional-scale scientific assessments, interviews with leading research groups conducting ongoing projects in the Carpathians, and a literature review using a key word search in Web of Science. The latter focused exclusively on peer-reviewed papers published in English language scientific journals.

The survey responses were coded to indicate the number of times particular risks, impacts and adaptation responses were mentioned. This was performed individually for each topic and then as a cross-cutting synthesis (or meta-analysis) across all the topics. Using the triangulation method, focal experts' shared risks and impacts of greatest concern were identified. When synthesized this way, survey responses were unequivocal with respect to the issues of central concern to national experts throughout the Carpathian region. The significance of these issues was subsequently validated by the literature review: the priority risks identified in survey results aligned closely with topics that are most actively investigated within recently published and ongoing forest science research.

JAN - JULY 2022

SURVEY ADMINISTERED TO FOCAL EXPERTS

2022 - 2023

LITERATURE REVIEW AND DATA PROCESSING

OCT 2023

ADOPTION OF THE ASSESSMENT AT THE 7TH MEETING OF THE CONFERENCE OF THE PARTIES TO THE CARPATHIAN CONVENTION

Table 3. Matrix structure from the survey for collecting key risks and adaptation responses for each priority topic.

PRIORITY TOPIC (e.g. Forest growth & productivity or Tree mortality)

Key risk s	Potential impacts (consequences, outcomes)

Adaptation Response Options for the Above Priority Topic

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)



Climate Change Risks and Impacts on Carpathian Forest Ecosystems and their Services



3.1 INTRODUCTION

Climate change poses direct risks to forest ecosystems by altering the competitive environment and causing physiological stress in constituent organisms (Figure 3). These direct risks are compounded by indirect risks caused by altered disturbance dynamics, such as windstorms, insect outbreaks and forest fires, particularly high-intensity events that remove forest canopies (Keeton et al. 2007).

Through the synthesis of survey responses (see Chapter 1.3), six key climate change related risks and impacts that pose significant threats to forest ecosystems, natural resources and human communities were identified. These risks represent potential vulnerabilities within the forest sector that necessitate adaptive planning and management (see Chapter 5).

The risks and impacts articulated by focal experts (survey respondents) closely mirror those identified by peer-reviewed scientific literature. These are also the subject of ongoing investigations at research institutions throughout the Carpathian region and Europe.





Figure 3. Climate change will affect Carpathian forests through both direct effects on the physiology, reproduction, and behavior of organisms and indirect effects on disturbance regimes that alter the competitive environment and, in some cases, increase exposure to the direct effects of climate change (e.g., temperature, precipitation, etc.). Figure modified from Keeton et al. (2007).

Figure 4. Chart depicting the process of identifying the risks and adaptation responses according to each priority topic.


TOPICS

Identified by the workshop at the Forum Carpathicum and expert subgroup meeting (see Chapter 2).

SEVEN PRIORITY TOPICS OF CONCERN



SURVEY QUESTIONS



SURVEY

Information requested in the survey administered to the focal experts for each of the 7 priority topics (see Chapter 2).

RESPONSES

Identified by the survey based on each of the 7 priority topics (see Chapters 3 and 4).

SIX PRIORITY RISKS

1. ALTERED DISTURBANCE REGIMES
2. ALTERED HYDROLOGIC REGIMES & FLOOD RISKS
3. INVASIVE SPECIES
4. CHANGES IN FOREST GROWTH & PRODUCTIVITY
5. ALTERED SPECIES COMPOSITION & DISTRIBUTION
6. FEEDBACK MECHANISMS & EFFECTS ON ECOSYSTEM SERVICES INCLUDING CARBON STORAGE

NINE ADAPTATION RESPONSES

1. ADAPTIVE SILVICULTURE
2. AFFORESTATION & REFORESTATION
3. CARBON FORESTRY
4. REWILDING & CORE AREA PROTECTION
5. INCREASE RESILIENCE TO DISTURBANCE
6. ASSISTED DISPERSAL & MIGRATION
7. MANAGEMENT PRACTICES TO MAINTAIN/IMPROVE THE ABILITY OF FORESTS TO RESIST PESTS & PATHOGENS
8. CLOSE-TO-NATURE SILVICULTURE/ECOSYSTEM-BASED MANAGEMENT
9. INTEGRATING FOREST-WATER INTERACTIONS INTO ADAPTIVE FOREST MANAGEMENT

3.2 SURVEY RESULTS AND LITERATURE REVIEW ON KEY RISKS AND IMPACTS

The six priority risks and impacts of concern are presented in the following section together with supporting information from the scientific literature:

1. Altered disturbance regimes
2. Altered hydrologic regimes & flood risks
3. Invasive species
4. Changes in forest growth & productivity
5. Altered species composition & distribution
6. Feedback mechanisms & effects on ecosystem services including carbon storage.

RISK: ALTERED DISTURBANCE REGIMES

The most frequently mentioned risk to all seven priority topics was the effects of climate change on disturbance regimes, both human and natural. Due to the land-use history of Carpathian countries, forests are already vulnerable to disturbances and studies project that climate change will increase their vulnerability to (1) pest and disease outbreaks, (2) windstorms, (3) drought and (4) wildfires.

Survey responses made clear the concern that increased disturbance risks will accelerate overall rates of forest change (see Risk: Altered species distribution and composition) and create feedback loops that diminish critical ecosystem services (see Risk: Feedback mechanisms and effects on ecosystem services including carbon storage).

The scientific literature also provides strong support for concern regarding disturbance risks expressed by focal experts, creating a consistent picture that disturbance events are becoming more frequent and severe in the Carpathians and that climate change is a contributing factor.



INCREASED PEST AND DISEASE OUTBREAKS

Warmer temperatures and altered precipitation patterns possibly increase pest and disease outbreaks. How these outbreaks manifest depend on the impacts of drought stress and high precipitation events:

- **DROUGHT STRESS:** Soil moisture deficits produce physiological stress that increases the susceptibility of trees to insect infestations. Warmer temperatures affect the reproductive cycles of bark beetles and other insect disturbance agents, potentially increasing the number of generations born within a single year (called “voltinism”) and leading to higher overall reproductive rates. Thus, bark beetle outbreaks in Carpathian spruce forests have been linked to warmer temperatures and drought

stress creating climate niches that increase both vulnerability of host trees and optimal pest reproductive success.

- **HIGH PRECIPITATION:** Warmer temperatures combined with sufficient rainfall and soil moisture create conditions that favor growth and dispersal of fungal tree diseases, such as *Armillaria* and *Phytophthora* root rots. These climate-related stresses will exacerbate the phenomenon of forest dieback, which is already severe in some areas, such as western Ukraine (Baradat et al. 2019). Additionally, compounded disturbance events may have cascading impacts on forest ecosystems, such as increased erosion, landslides, and soil compaction (Hlásny et al. 2017).

INCREASED DROUGHT

The second most frequently mentioned risk was drought. Recent climate trends and severe drought events suggest that drought and associated disturbance risks are increasingly frequent and severe within the Carpathian region and Europe as a whole (Science for Environment Policy 2021). Focal experts identified that drought poses grave consequences for forest growth and productivity, regional tree mortality rates, biodiversity, future shifts in species composition and the climate regulating functions of Carpathian forests.

The connection to forest-derived water resources and other ecosystem services, such as carbon storage, is also clear. There is great concern that droughts will interact with other stresses, such as wind, fire, insects and pathogens, to reduce tree growth rates, increase mortality, alter tree regeneration dynamics, and exacerbate forest decline, not just in spruce forests but other types as well, including beech (Petritan et al. 2015; Thom et al. 2023). The persistent and sometimes lagged (or multi-year) influence of droughts on disturbance risks is also likely to have major structuring effects on temperate forests into the future (Senf & Seidl 2021b).



INCREASED WILDFIRES

Forest fire risks are highly likely to increase with greater interannual variability in seasonal precipitation, which lengthens dry seasons and increases drought frequency, warmer temperatures and heat waves. These climatic changes will directly influence key drivers of fire ignition and spread, including ambient relative humidity and fuel moisture content, thereby exacerbating forest fire risks.

Greater fire activity is likely to increase rates of vegetation transition between plant community types, as fires disturb forest canopies and alter the regeneration environment. The resulting shifts in forest composition and structure may have pros and cons: helping to diversify landscapes and increasing carbon uptake rates in successional stands, but increasing declines in desirable forest types, shifting landscape to young seral stages, reducing net carbon storage, and exacerbating spread of invasive species (Costa et al. 2020).

INCREASED WINDSTORMS

Increased frequency and intensity of windstorms result in greater tree mortality and associated changes in the production of critical ecosystem services, including habitat, hydrologic regulation, carbon storage, and timber resources (Seidl et al. 2014; Grodzki et al. 2018). The structure, composition, and age class distribution of Carpathian forests are likely to shift in response to an increased prevalence of severe windstorms. Wind disturbances can also interact with other disturbances by increasing the availability of dead and dying host trees for insects and pathogens and enhanced fuel loading as a major fire risk.



CARPATHIAN AREAS OF CONCERN

Drought stress is a major concern for forests in the Carpathians, particularly in the southern and eastern parts of the range where soil moisture availability is more limited (Hlásny et al. 2017). Fire frequencies have already been shown to be increasing in portions of the eastern and southern Carpathians (Kelley et al. 2019) and relative to Europe overall, the Carpathian Mountains have been assessed to have particularly high vulnerability to climate change-driven forest fires (Forzieri et al. 2020).

There is also clear evidence that drought risks will be more pronounced in some sub-regions and locales than others, such as particular sub-basins in the Polish Tatra Mountains (Bokwa et al. 2021).

It is also important to note that disturbances play a critical structuring role in Carpathian forest ecosystems. When operating within a more natural range of variability (i.e. one less altered by climate change and historically poor forest management), disturbances (1) create biodiverse habitats and (2) maintain critical carbon processes.

Forest stands that regenerate after experiencing disturbances produce diverse successional habitats and stand structural elements that many species depend on. The formation and persistence of species diversity and critical habitats, such as “natural” and virgin forests and old forest elements like large legacy trees within younger stands depend on natural disturbance and successional processes. Many of these stand structural conditions buffer microclimates beneath complex forest canopies and maintain high

genetic diversity, fostering climate change resilience through evolutionary potential. Additionally, disturbances maintain carbon environments and processes, including forest patches with high levels of carbon uptake (sequestration) and patches with high levels of biomass (storage) (Mikolas et al. 2021, 2023). A key element of patch mosaics is the inclusion of the full diversity of seral (i.e. successional) habitats and stand structural conditions upon which many species depend, while also maintaining an optimal mix of carbon flux pathways and pools.

Thus, restoring the forest conditions needed to support desired disturbance activity levels may be one of many adaptation approaches (Aszalos et al. 2022). However, while some survey responses emphasized these functions, many others focused solely

on the threats, illustrating differing viewpoints around this issue.

The identified risks align closely with the leading topics in current forest science research in Europe. Both national experts and scientific literature suggest various adaptation responses, including forest management to increase compositional heterogeneity across landscapes, restoration of mixed-species and beech forests where these were historically endemic, and management for forest structures that are less susceptible to disturbances (see Chapter 4). Recent assessments also stress the importance of not just protecting forests from disturbances, but rather restoring diverse landscapes that are more resilient through natural recovery processes and adaptive silvicultural practices (Lindner et al. 2020).

RISK: ALTERED HYDROLOGIC REGIMES AND FLOOD IMPACTS



ALTERED HYDROLOGIC REGIMES

Focal experts identified altered hydrologic regimes to be a major vulnerability within the region and one that is caused by altered disturbance risks and human impairment of watershed functioning. Regarding the former, respondents stressed that large-scale disturbances, such as fire, bark beetle outbreaks and defoliating insects, will reduce water uptake by trees and reduce infiltration into soils. Regarding the latter, poorly designed forest roads, development, and impervious surfaces change the form of the hydrograph, meaning that water after precipitation events is delivered more rapidly and at greater volumes directly to streams, rivers, and other surface waters. Collectively, these interacting climate and human impacts increase runoff and the


intensity of peak flows, thereby inducing severe erosion, flooding during high precipitation events, and possibly chemical loading.

FLOOD IMPACTS

Focal experts identified flood risks and their connection to forest cover and management as a major risk of concern. Concern over floods risks may seem idiosyncratic given an overall trend of increasing drought frequency, but there was acute awareness of vulnerability to extreme precipitation events and the likelihood that the Carpathians, like much of Europe, will experience more of these in the future.

RISK: INVASIVE SPECIES

Warmer temperatures and altered precipitation patterns may increase risks and vulnerabilities associated with the spread of invasive insect pests, tree pathogens, and noxious plants, such as Japanese knotweed and mistletoe.



RISK: CHANGES IN FOREST GROWTH AND PRODUCTIVITY

Survey responses showed a clear concern for the impacts of climate change on forest growth and productivity, including the potential for declines, but acknowledged that these impacts will be expressed differently throughout the region. According to focal experts, productivity responses will vary by climate sub-region, topographic position, and plant species. These variations depend on climate change's direct effects on tree physiology, phenology, and reproductive success and its indirect effects on the dynamics of natural disturbance regimes.

For example, warming trends from 1987 to 2022 across Romania resulted in "greening" (increased productivity) in higher elevation forests, while some lowland forests showed a degree of "browning" (decreased productivity), based on assessments using remotely-sensed Normalized Difference Vegetation Index (Prăvălie et al. 2022).

ALTERED PHENOLOGY

Increased temperature and variations in precipitation were the most cited drivers of productivity impacts by focal experts, which the scientific literature also explores across the Carpathians. Warmer spring temperatures can lead to earlier budburst and production of foliage (i.e. altered phenology), which can influence a variety of ecosystem processes, including the availability of sunlit patches for spring ephemerals, nutrient uptake rates, and tree growth associated with growing season lengths (Vitasse et al. 2018; Mihai et al. 2018). With a longer growing season, increased water uptake by trees and other plants may lead to soil moisture deficits if precipitation is also lower (Buras et al. 2018), creating negative feedback on forest productivity (Dobrowolska et al. 2017; Kruhlov et al. 2018).

Other effects associated with altered phenology include the timing of insect emergence and bird migration (Buras et al. 2016), which pose risks for biodiversity if other environmental conditions, like populations of natural

predators or food availability, do not shift congruently. And where phenology does not keep up with the rate of warming, tree species may become less competitive and more prone to niche displacement or invasion by non-native species.

ALTERED CO₂ FERTILIZATION

Respondents' views differed on the potential for CO₂ fertilization—the phenomenon of a CO₂-enriched environment producing higher photosynthesis efficiency in plants—to enhance forest productivity. Respondents were aware of conflicting scientific evidence on this topic: it depends on the time frame of analysis, interactions with other stressors like moisture availability, and nutrient limitations that can lead to decreased productivity over time (Penuelas et al. 2020).

RISK: ALTERED SPECIES COMPOSITION AND DISTRIBUTION

Focal experts from all Carpathian countries expressed concern regarding the future of their natural heritage. While not mentioned as frequently as disturbance or drought risks, the effects of climate change on species and biological diversity were highlighted as a major threat in every focal expert response. Additionally, increased disturbance risks will accelerate overall rates of forest change, exacerbating other threats such as the spread of invasive species, species range shifts, and loss of important habitats for biodiversity.

Of particular concern are species with intrinsically low dispersal rates and species in isolated habitats. These species face climate-related extinction risk, especially those on mountain tops and in highly fragmented landscapes. Many respondents stressed the

likely impacts of climate change on keystone and flagship species and the overall composition of forest communities. Climatic factors and anthropogenic pressures interact to shift habitats, a fundamental risk to the viability of at-risk populations of plants, wildlife, and other taxa. There was consensus that these risks require broader, coordinated adaptation responses to minimize the potential for local or broad scale extirpations and to facilitate species range shifts.

The survey responses are supported by scientific literature, which predicts that Carpathian tree species will respond individually to the direct effects of climate change. Risk factors vary by species according to the different physiological mechanisms linking climate to growth and survival, such

as hydraulic failure, frost damage, and carbon starvation (Petit-Cailleux et al. 2021). Some tree species are likely to experience increased growth and competitiveness, while others will become less competitive or face declined habitat availability (Kobiv 2018). In particular, non-native species possessing future-adapted traits may become more competitive, allowing them to become more widely invasive (Mátyás et al. 2019). According to the literature, it remains uncertain whether species ranges will be able to adapt to rates of climatic change and migrate to and colonize new locations within their environmental tolerance ranges (Thuiller et al. 2005). Higher tree mortality is predicted to increase rates of succession and compositional change across European forests (Ruiz-Benito et al. 2017).



As temperature and precipitation patterns change across the Carpathian Mountains, the competitive environment will be altered, favoring some tree species over others depending also on the site characteristics and local growing conditions. For example:

- Less drought tolerant tree species at lower elevations may expand their ranges to higher elevations where moisture availability is higher, resulting in elevational shifts in the relative abundance of tree species (Grodzki et al. 2018). In boreal forest types, montane spruce-fir and deciduous species like oak, beech and maple are likely to increase in relative abundance. At higher elevations, the composition of subalpine forests will be increasingly dominated by mesic species such as spruce, fir, and beech.
- Drought tolerant species, such as oak, pine, and fir, are predicted to increase in relative dominance, particularly on drier sites, as species requiring higher soil moisture, such as beech and spruce, decline. Researchers predict that beech-dominated forests may decline by 50% by the 22nd century and spruce forest areas will shrink by 60%, whereas oak forests are likely to increase by 40% in the Carpathians (Hlásny et al. 2017; García-Duro 2021).




Table 4. Responses to climate change effects for selected Carpathian tree species. The table synthesizes information compiled by Kholiavchuk, Gurgiser, and Mayr (2023) through a comprehensive Web of Science search for literature addressing climate change effects on Carpathian tree species. Data from that study is supplemented with information obtained through this assessment.

SPECIES	GROWTH RESPONSES	RESPONSE TO DROUGHT	PREDICTED RANGE SHIFTS
NORWAY SPRUCE (<i>Picea abies</i>)	<ul style="list-style-type: none"> • Increased variability of radial growth at lower elevations • May be more adaptive to increased temperatures at higher elevations 	<ul style="list-style-type: none"> • Reduced growth and survival in response to extreme droughts • Synergistic interactions with bark beetle susceptibility causing declines 	<ul style="list-style-type: none"> • Climate change contributes to encroachment of spruce into alpine meadows, although the stronger influence is pastureland abandonment • Predicted increase in dominance in sub-alpine zones • Predicted overall decrease in abundance
BEECH (<i>Fagus sylvatica</i>)	<ul style="list-style-type: none"> • Growth and survival more strongly affected by drought than by heat 	<ul style="list-style-type: none"> • Highly susceptible to drought during the mid to late growing season 	<ul style="list-style-type: none"> • Declines on drought-prone sites • Range expansion at higher elevation on mesic sites and within the boreal zone • Predicted overall decrease in abundance
SILVER FIR (<i>Abies alba</i>)	<ul style="list-style-type: none"> • Growth may increase with higher temperatures in some regions 	<ul style="list-style-type: none"> • Variable responses • Relatively adaptive on drier sites 	<ul style="list-style-type: none"> • Predicted range expansion on drier sites
OAK SPECIES (<i>Quercus</i> spp.)	<ul style="list-style-type: none"> • Adaptive to temperature increases 	<ul style="list-style-type: none"> • Exhibits greater drought resistance compared to other Carpathian tree species • Severe drought increases risk of fungal and mistletoe infection 	<ul style="list-style-type: none"> • Range expansion where oak becomes more competitive, particularly in the southern Carpathians • Predicted overall increase in abundance

RISK: FEEDBACK MECHANISMS AND EFFECTS ON ECOSYSTEM SERVICES INCLUDING CARBON STORAGE

Finally, a major theme in survey responses was the potential for climate change to alter the mix of ecosystem services provided by Carpathian forests through a variety of feedback mechanisms. Based on survey responses and supporting literature, this assessment finds that complex interactions between climate change and forest ecosystem processes, such as productivity, carbon, and water cycles, may create feedback loops that accelerate rates of change both in the climate system and in forest resources.

CARBON FEEDBACKS

Focal experts were particularly concerned about greater rates of carbon flux to the atmosphere due to disturbances, drought stress, and reduced forest productivity, reducing the effectiveness of Carpathian forests as a NCS.

Increased net carbon fluxes to the atmosphere could potentially exacerbate both regional and global climate change. Carpathian forests store approximately 6% of Europe's total forest carbon stock (Zlatanov et al. 2016). However, the direct and indirect effects of climate change are likely to lead to a significant decline and possibly a reversal in sink capacity (Kruhlov et al. 2018). In a 500-year projection of climate change impacts on the Ukrainian Carpathians, Kruhlov et al. (2018) found that, due to the combined effects of compositional changes and disturbances, aboveground carbon storage in forests declined between 6.6% to 20.6% depending on the climate scenario.

These results were supported by Pfeifer et al. (2018), who predicted that the Carpathian Mountains will become a carbon source over the current century due to the drivers identified by Kruhlov et al. (2018).

DISTURBANCE FEEDBACKS

Disturbance impacts will create feedback loops that diminish the provisioning of critical ecosystem services, including timber and non-timber resource production, carbon storage, and hydrologic regulation.

Forests' increased vulnerability to disturbances described earlier will pose significant management challenges from damage to timber resources and the need to manage for resilient landscapes (Hlásny et al. 2019). For example, positive feedbacks link climatic conditions and the water cycle to the distribution of forest types, their age class distributions, stand structure characteristics influencing hydrological regulation, and their associated biomass and

carbon storage (Ceballos et al. 2018; Kruhlov et al. 2018). There are critical interactions between disturbance types, increasing system vulnerability overall (Thom & Seidl 2016; Seidl et al. 2017).

Carpathian forests have heightened vulnerability to climate risks due to historic forest management and land-use changes which have homogenized forest composition and structure. These have increased susceptibility to disturbance and forest health risks, creating positive feedback loops that are likely to accelerate change (Keeton et al. 2013; Hlásny et al. 2017).

DISTURBANCE IMPACTS ON CARBON STORAGE

The literature illustrates that forest carbon storage decreases as disturbance frequencies increase (Seidl et al. 2014). The temporal dynamics of the carbon balance following very large disturbances can be highly complex, as shown after a major wind event that affected more than 12,000 ha of mature forest in the Tatra Mountains in 2004 (Fleischer et al. 2020). Immediately after the event, forests shifted to carbon sources, only later becoming carbon sinks with the growth of successional vegetation. Areas infested with bark beetles after the wind event remained carbon sources for longer periods (Fleischer et al. 2020).

On the other hand, landscapes that are diversified to enhance resilience may accommodate a range of disturbances closer to historic ranges of variability and a more optimal mix of carbon sequestration and storage and seral habitats for biodiversity (Mikolas et al. 2021; Aszalos et al. 2022).



HYDROLOGIC FEEDBACKS

Feedbacks between climate change and water resources posed another great concern to the focal experts.

Hydrologic regimes may be especially sensitive to shifts in seasonal precipitation and evapotranspirative demand, leading to changes in both the quantity and quality of freshwater resources. These effects will be distributed unevenly both spatially and seasonally, with annual precipitation predicted to increase in the northern Carpathians and decrease in the southern portion of the region (Werner et al. 2016).

Research suggests that stream flow, soil moisture availability, and clean water production will decline the most during the summer months and in the drier south, while 30-year flood risks will increase slightly to strongly, although predictions are sensitive to uncertainties in climate scenarios and data limitations (Didovets et al. 2019). Feedbacks between climate change and forest productivity (negative reinforcement) and disturbances like forest fires (positive reinforcement) are highly probable, with significant implications for regional economies, livelihoods, and human communities.

ALBEDO FEEDBACKS

An often poorly recognized yet critically important feedback between climate and forests relates to albedo, or reflectivity of the Earth's surface. Vegetation types and land surfaces with lower albedo reflect less of the incoming shortwave (or ultraviolet) radiation, converting more of it to long wave radiation (or heat). Forest ecosystems typically have lower albedo and thus absorb more solar radiation compared to other land cover types, such as grasslands or agriculture (Drever et al. 2021). If oak and pine forests increase in relative abundance in the Carpathians as predicted, this is likely to decrease the region's albedo, since beech and spruce forests typically have higher albedo (Vincze et al. 2019). This would create a positive feedback loop, exacerbating a warming regional climate.

PRODUCTIVITY AND RELATED ECONOMIC FEEDBACKS

As identified by both focal experts and the literature, climate change is expected to lead to major vulnerabilities related to forest ecosystem growth and productivity in the Carpathians. For example, beech forests in the Polish Carpathians are predicted to experience a 40% decrease in productivity by the year 2100 in response to rising temperatures and declining summer rainfall (Czajkowski et al. 2017). Such productivity declines would have significant negative economic effects, feeding back on forest sector revenue and ecosystem services underpinning the sustainable management of these systems.



3.3 AREAS OF CONTINUING DEBATE

Overall, the risks and impacts articulated by focal experts closely mirror those identified by peer-reviewed scientific literature. However, respondents had different views on some issues, such as restoration of older forests, the carbon sequestration and storage value of older forests, and whether forest management intensity should be increased or decreased.

Respondent views sometimes diverged from the developing consensus within the scientific literature, for example on the carbon value of older forests (Keeton et al. 2010; Mikolas et al. 2023). However, the literature review showed the same degree of debate on the topic of optimal forest management intensity, suggesting that the focal experts are not alone in having reached widely different conclusions. There was general support for forest restoration, afforestation and reforestation, and conversion cutting to restore endemic species composition and to create more heterogeneous landscapes.

Table 5. Correlation of risks identified by focal experts according to each of the seven priority topics.

PRIORITY TOPICS	RISKS OF CONCERN					
	ALTERED DISTURBANCE REGIMES	ALTERED HYDROLOGIC REGIMES & FLOOD RISKS	INVASIVE SPECIES	CHANGES IN FOREST GROWTH & PRODUCTIVITY	ALTERED SPECIES COMPOSITION & DISTRIBUTION	FEEDBACK MECHANISMS & EFFECTS ON ECOSYSTEM SERVICES, <i>Including Carbon Storage</i>
FOREST GROWTH & PRODUCTIVITY	Correlated	Correlated	Not Correlated	Correlated	Not Correlated	Correlated
BIOMASS & CARBON STOCKS	Correlated	Not Correlated	Not Correlated	Correlated	Not Correlated	Correlated
TREE MORTALITY	Correlated	Not Correlated	Not Correlated	Not Correlated	Not Correlated	Correlated
CHANGES IN SPECIES RANGE & ABUNDANCE AND HABITAT SHIFTS	Correlated	Not Correlated	Not Correlated	Correlated	Correlated	Not Correlated
INVASION BY NON-NATIVE SPECIES	Correlated	Not Correlated	Correlated	Correlated	Correlated	Not Correlated
FOREST ECOSYSTEM SERVICES	Correlated	Correlated	Not Correlated	Not Correlated	Not Correlated	Correlated
FOREST-WATER INTERACTIONS, <i>Including Hydrologic Regulation & Riparian Dynamics</i>	Correlated	Correlated	Not Correlated	Correlated	Not Correlated	Correlated



4 Adaptation Approaches

4.1 INTRODUCTION

This chapter aims to elaborate on adaptation approaches for forests and the forest sector in the entire region based on survey responses. Focal experts were asked to indicate adaptation response options for addressing each priority topic and briefly highlight their intended effects, and identified the following adaptation responses:

1. Adaptive silviculture
2. Afforestation & reforestation
3. Carbon forestry
4. Rewilding & core area protection
5. Increase resilience to disturbance
6. Assisted dispersal & migration
7. Management practices to maintain/improve the ability of forests to resist pests & pathogens
8. Close-to-nature silviculture/ecosystem-based management
9. Integrating forest-water interactions into adaptive forest management

The most frequently mentioned adaptation response was forest restoration, while sustainable management—including broader use of close-to-nature silviculture and continuous cover forestry—was the second-most mentioned response.

4.2 RESPONSE OPTIONS ALONG IDENTIFIED KEY TOPICS

The following pages provide a synthesis of responses received for each of the key topics with support from related literature.



ADAPTATION RESPONSE:

ADAPTIVE SILVICULTURE

CHARACTERISTICS:

Emphasizes ecological principles and aims to maintain or enhance the natural processes and functions of the forest ecosystem, such as biodiversity, soil health, water quality, and habitat connectivity.

Balances multiple objectives, such as timber production, wildlife habitat conservation, water balance and carbon sequestration, disturbance processes and recreational opportunities, based on the specific context and goals of forest management.

Harnesses both natural and artificial regeneration to shift or convert species composition to mixed forests where these were historically endemic or where they will be future adapted. This may take the form of a variety of silvicultural approaches, including but not limited to close-to-nature forest management, which is a set of practices to confer resistance or resilience to climate change:

- Continuous cover forestry with uneven aged, diverse forests
- Stocking management, including thinning to reduce stand densities
- Diversified landscape mosaics in terms of patch structure and composition
- Gap- and retention-based regeneration harvesting systems
- Use of prescribed burning in forest types and drier sites that once supported low intensity, ground fires

Requires ongoing monitoring of forest conditions and response to management actions. This helps in assessing the effectiveness of different approaches and making informed decisions.

KEY ADVANTAGES:

- Creates greater resilience to abiotic damage in comparison with even-aged stands.
- Improves biodiversity through the creation of a vertically and horizontally diverse habitats at stand scales and mosaics of seral habitats at landscape scales.
- Leads to more diverse landscape that limit disturbance spread and optimize ecosystem services such as carbon sequestration and storage.

PRIORITY TOPIC ADDRESSED:

1 FOREST GROWTH & PRODUCTIVITY

MAIN RISKS ADDRESSED:

- Spruce mortality and decline in vigor of other forest types, including beech
- Drought and increased disturbance risks, such as bark beetles, wind, and forest fires
- Increased physiological stress in trees leading to reduced growth and vigor
- Landscape-scale continuity of vulnerable host trees for insect pests

INTENDED EFFECTS:

Better adaptation to future climatic conditions and resilience or resistance to a variety of stresses.

Decreased drought- and disturbance-related risks.

Reduced wildfire risk.

Managing tree density and thinning removes shrubs and flammable vegetation in between the ground and the crown level (vertical continuity). Species with a higher age mix and species mix tend to be less vulnerable to wildfires compared to mono-age and mono-species stands, since their more complex structure can slow fire spread. Restores capacity for low intensity fires that do not "canopy out" and become high intensity, stand replacing burns.

Rehabilitated and sustainable ecosystems.

Continuation of native tree cover having usually better productivity than invasive species.

Increased functional and structural diversity.

POTENTIAL DISADVANTAGES:

- Time constraints and costs in achieving irregular structures through lost production during the transformation period.
- Lack of knowledge of the process of transformation.
- Limited range of sites where transformations may be possible.

ADAPTATION RESPONSE:

AFFORESTATION & REFORESTATION

CHARACTERISTICS:

Afforestation: Converting land that has not been tree-covered for at least 50 years (according to UNFCCC) into forest.

- Establishment of forests in areas that historically have not been forested, such as barren lands, agricultural fields, or urban areas. It involves planting tree seedlings or direct seeding, and sometimes soil preparation and protection from herbivores and competing vegetation, to initiate the growth of new forests.
- Conversion of non-forest land, for example, through land-use planning, land acquisition, and implementation of suitable silvicultural practices.
- Careful consideration is needed for selecting appropriate tree species that are well-suited to the local and future climate and soil conditions and intended objectives. The selection may include native species, non-native species, or a combination of both, depending on the goals, ecological context and applicable legal frameworks. Use of climate forecasts to estimate future site-specific growing conditions is advised.

POTENTIAL DISADVANTAGE:

- High initial investment to establish new stands coupled with the several-decade delay and the need for subsequent pre-commercial thinning until afforested areas generate revenue or provide desired ecological functions. This may represent a major constraint.

PRIORITY TOPIC ADDRESSED:

1 FOREST GROWTH & PRODUCTIVITY

MAIN RISKS ADDRESSED:

- Floods and landslides
- Droughts, extreme temperatures, water scarcity
- Soil degradation
- Biodiversity loss
- Declines in rural livelihood opportunities

Reforestation: Converting recently non-forested land into forest.

- Focuses on restoring forest cover in areas that have been deforested or significantly degraded. It involves the replanting or natural regeneration of trees in areas that were previously forested but have experienced forest loss due to clearing, logging, fire, or other disturbances.
- Aims to restore ecological integrity and functionality to degraded forest ecosystems. It involves the reintroduction of native tree species and other associated vegetation, promoting natural processes, and recovering biodiversity and ecosystem services.
- Often requires site preparation activities such as removing debris, controlling competing vegetation, and improving soil conditions to facilitate successful tree establishment. Careful management practices, including monitoring and maintenance, are implemented to ensure the long-term success of the restored forests.
- Often considers appropriate game management and landscape-scale planning to enhance connectivity between fragmented forest patches. This helps to support wildlife movement, genetic exchange, and ecosystem resilience.
- Should consider socio-economic factors, such as local community involvement, livelihood improvement, and sustainable forest management practices. It can provide economic opportunities through sustainable timber production, non-timber forest products, and ecotourism. Gap-based and retention-based regeneration harvesting systems.
- Use of prescribed burning in forest types and drier sites that once supported low intensity, ground fires.

INTENDED EFFECTS:

Enhanced resilience to disturbances.

Landscape diversification increases the complexity and diversity of patch mosaics to limit disturbance contagion and spread.

Reduced destruction or degradation of habitats and safeguarded ecosystem services.

Reduced fragmentation/increasing ecological connectivity.

Facilitates species migration under climate change conditions.

Reduced risk of landslides, restored riparian and floodplain functionality, protected water bodies from sedimentation, and promoted healthy watershed management.

Replenished groundwater, reduced runoff, and improved water availability and quality.


Provision of valuable resources for local communities, such as timber, non-timber forest products, wood bioenergy, and medicinal plants.

A renewable source of timber and other forest products supports the forestry industry and contributes to sustainable rural economic development.

Increased local community involvement.

Fosters a sense of ownership, empowerment, and stewardship. It can strengthen community resilience, promote social cohesion, and enhance cultural values associated with forests.

Increased opportunities for forest-based recreation, including hiking, biking, skiing, wildlife observation, and nature tourism.



ADAPTATION RESPONSE:

CARBON FORESTRY

CHARACTERISTICS:

Planting or natural regeneration of future-adapted tree species, producing diverse forests and landscape mosaics with high carbon sequestration and storage capacity.

Using carbon forestry practices that yield high biomass/high carbon storage forests over time. These include stand improvement thinning, extended harvest rotations, retention systems for regeneration harvests, close to nature silviculture, and unmanaged inclusions or reserves. Support chosen management practices for offsetting carbon emissions.

Combining these practices to mitigate climate change by capturing carbon dioxide (CO₂) from the atmosphere and storing the carbon in forest biomass, both above and belowground, thereby offsetting GHG emissions from other sources.

KEY ADVANTAGES:

- Increases carbon sink capacity and stocks, substitution effects when wood products replace materials with high carbon footprints, such as plastic, steel, and concrete.
- Generates carbon credits, also known as forest carbon offsets, which can be sold in carbon markets or used to offset emissions from other sectors.
- Carries co-benefits like provision of complex, late-successional habitats for biodiversity, riparian functionality, and hydrologic regulation.

POTENTIAL DISADVANTAGES:

- Some carbon forestry practices, such as extended rotations, carry heightening disturbance risks and therefore must be planned carefully at landscape scales.
- There can be tradeoffs with some forms of timber production and is therefore best planned holistically alongside other forest management objectives.

PRIORITY TOPIC ADDRESSED:

2 BIOMASS & CARBON STOCKS

MAIN RISKS ADDRESSED:

- Forest management intensity and disturbance frequency
- Changing/reduced carbon uptake and carbon dynamics (sequestration, storage, and fluxes)

INTENDED EFFECTS:

Maintained and/or enhanced stored carbon.

Carbon stored in terrestrial ecosystems is vulnerable to greater flux rates (i.e. CO₂ flux to the atmosphere) as the climate changes and is dependent on the specific mix of economic uses of forests, including conversion to non-forest land-uses. Sustainably managed, protected forests act as carbon sinks and storage reservoirs.

Carbon is stored belowground in the soil system and in the forest long-term, as well as in durable (long-lasting) wood products.

Enhanced removal of CO₂ from the atmosphere (carbon sequestration), thereby helping to mitigate climate change.

Increased wood processing efficiency, re-use, and recycling.

The carbon benefits depend on the overall portfolio of forest carbon management approaches and their net effect.

Generated emissions offsets, which are tradable credits representing the additional reduction or removal of greenhouse gas emissions over a baseline.

These offsets can be sold or used to compensate for emissions in other sectors, such as energy production or transportation, effectively neutralizing a portion of their carbon footprint.

Socio-economic benefits brought to local communities.

Jobs created in tree planting and forest management, income generated from carbon credit sales, and opportunities created for sustainable land use practices.

ADAPTATION RESPONSE:

REWILDING & CORE AREA PROTECTION

PRIORITY TOPIC ADDRESSED:

2 BIOMASS & CARBON STOCKS

MAIN RISKS ADDRESSED:

- Changing/reduced carbon uptake and carbon dynamics (sequestration, storage, and fluxes)

CHARACTERISTICS:

Rewilding: Conservation efforts aimed at restoring and protecting natural ecosystems processes and wilderness areas that will involve fewer active forms of natural resource management.

- May include recreation management, invasive species control, use of prescribed fire, and other reserve-based management practices.
- Reintroducing species that have become locally extinct or have declined due to human activities. These species are typically keystone species or ecosystem engineers that play critical roles in shaping their habitats. By reintroducing such species, ecosystem services can be restored, and habitats can be revitalized.
- Reinstating ecological processes, including predation, herbivory, natural disturbance dynamics, and ecological succession, which can have cascading effects throughout the ecosystem.

Core area protection: Designating and safeguarding specific areas within a larger landscape or ecosystem for the conservation of biodiversity and ecological processes.

- Often involves the establishment of protected areas, such as national parks, nature reserves, wildlife sanctuaries, or other forms of protected land.
- Formation of contiguous areas of natural forest linked by forests and landscapes managed for ecological connectivity.

INTENDED EFFECTS:


Restored overall health and functionality of entire ecosystems.

Revitalise natural processes by rewilding as a Nature-based Solution (NbS).

High levels of carbon storage.

Core reserves complement more intensive forest management employed elsewhere on the landscape, thereby providing "risk spreading".

May result in reduced net carbon uptake over centuries but compensates with high levels of carbon storage and continued high rates of carbon sequestration over the near to mid-term as forests recover and develop towards a late-successional condition.





KEY ADVANTAGES:

- **Ecosystem Restoration:** Rewilding can restore ecological processes and functions that have been disrupted due to human activities. This includes natural predator-prey dynamics, seed dispersal, pollination, vegetation succession, and nutrient cycling. Restoring these processes can have cascading positive effects on the entire ecosystem.
- **Core area protection** is one of the most effective ways to conserve biodiversity and protect sensitive ecosystems. By establishing protected areas, critical habitats can be preserved, allowing for the conservation of endangered species, rare plants, unique ecosystems and their ecosystem services.
- **Carbon sequestration:** Rewilded areas often have increased vegetation cover and a greater variety of plant species, which can enhance carbon sequestration. This helps mitigate climate change by reducing atmospheric carbon dioxide levels and storing carbon in soils and vegetation.
- **Ecotourism and Economic Benefits:** Rewilded areas can attract tourists and nature enthusiasts, creating economic opportunities for local communities. Ecotourism related to rewilding projects can generate revenue, job opportunities, and support local businesses, contributing to sustainable development.
- **Scientific and Educational Value:** Core areas offer opportunities for scientific research and education. They provide undisturbed ecosystems for studying natural processes, ecological interactions, and species behavior. These areas also serve as outdoor classrooms and living laboratories, contributing to ecological knowledge and environmental education.

POTENTIAL DISADVANTAGES:

- **Human-Wildlife Conflicts:** Introducing or reintroducing large predators or other wildlife species can lead to conflicts with human activities such as agriculture, livestock farming, and infrastructure development. Predation on livestock and crop damage may create challenges and tensions between local communities and rewilding initiatives. On the other hand, there are many examples of community-based partnerships formed to overcome these challenges, leading to successful reintroduction programs.
- **Change in the type and distribution of flora and fauna:** Introducing new species or increasing the cover of historical endemic vegetation can shift the landscape from current or cultural norms, sometimes requiring an adjustment in cultural aesthetics, uses, and acceptance.
- **Uncertainty and Long Timeframes:** Rewilding is a complex process that requires careful planning and implementation. It can take many years or even decades for ecosystems to fully recover and for rewilding efforts to demonstrate their intended benefits. There is inherent uncertainty in predicting outcomes and the success of rewilding initiatives particularly in the face of climate change and altered successional dynamics.
- **Restricted Land Use:** The establishment of core areas often involves restricting or prohibiting certain land uses and activities. This can create conflicts with local communities, landowners, and resource-dependent industries such as agriculture, logging, or mining. Balancing conservation goals with socioeconomic needs and ensuring equitable solutions is a challenge.

ADAPTATION RESPONSE:

INCREASE RESILIENCE TO DISTURBANCE

CHARACTERISTICS:

Enhances and maintains species, structural and genetic diversity by favoring existing genotypes that are better adapted to future conditions, incorporating genetic diversity from a greater range of population sources, and including pest- or drought-resistant varieties where appropriate.

More intensified thinning practices and care of forest stand edges.

Promotes redundancy of ecological representation within core protected areas. Also “functional redundancy” which means having multiple species or ecological components that perform similar functions, providing compensatory capacity if one species declines or is adversely affected by climate change. This functional diversity ensures that multiple ecological processes and services are maintained, even if some species or functional groups are lost or impacted.

Establishes ecological corridors and maintains landscape connectivity to facilitate the species’ range shifts, dispersal and genetic interchange among populations, and continuation of ecological processes. Connected landscapes allow for the dispersal of species, enabling recolonization and gene flow following disturbances. Corridors can also help species adapt to shifting environmental conditions caused by climate change.

KEY ADVANTAGES & POTENTIAL DISADVANTAGES:

Depends on approaches to increase resilience.

PRIORITY TOPIC ADDRESSED:

3 TREE MORTALITY

MAIN RISKS ADDRESSED:

- Increasing soil moisture deficits
- Prolonged drought due to reduced precipitation and higher temperatures likely in some areas

INTENDED EFFECTS:

Enhanced diversity in forests.


Diverse forests exhibit higher variability in resistance to pests, drought and access heat.

Lowered competition and the probability of drought-related tree mortality.

Involves reducing stand densities, for instance in intensively managed coniferous forests.

Enhanced complexity and diversity of patch mosaics (e.g., different types and ages of vegetative communities) across the landscape.

Helps limit contagion and spread of insects and plant diseases.



ADAPTATION RESPONSE:

ASSISTED DISPERSAL & MIGRATION

CHARACTERISTICS:

Moving species or genotypes to new locations that should better match their climatic suitability in the future through:

- Assisted population migration (also assisted genetic migration or assisted gene flow): moving seed sources or populations to new locations within the historical species range; or
- Assisted range expansion: moving seed sources or populations from their current range to suitable areas just beyond the historical species range (adjacent areas), facilitating or mimicking natural dispersal; or
- Assisted species migration (also species rescue, managed relocation, or assisted long-distance migration): moving seed sources or populations to a location far outside the historical species range, beyond locations they would naturally spread.

INTENDED EFFECTS:

Maintained forest productivity.

Particularly when suitable seed sources or populations are moved to locations within, or just beyond, the historical range where growing conditions are likely favorable under future climatic conditions.

Maintained crucial ecosystem functions like wildlife habitat, carbon sequestration, and watershed services under ecosystem restoration conditions.

Control of herbivory on tree seedlings is a critical component of this approach.

Populations and species successfully moved across ecological barriers in fragmented landscapes, such as roads, agriculture, cities, and other human infrastructure.

PRIORITY TOPIC ADDRESSED:

CHANGES IN SPECIES RANGE AND ABUNDANCE & HABITAT SHIFTS

MAIN RISKS ADDRESSED:

- Extinction risk for species with intrinsically low dispersal rate and species in isolated habitats such as mountain tops
- Impacts on keystone and flagship species and the composition of forest communities
- Habitat shifts through interaction of climatic factors and anthropogenic pressures





KEY ADVANTAGES:

- **Ecosystem Restoration:** Especially in areas where species have been extirpated or ecosystems have been degraded. It can aid in the reestablishment of natural ecological communities and the recovery and preservation of ecosystem processes such as such as pollination, seed dispersal, and nutrient cycling.
- **Conservation of Species:** Especially species that are at risk of extinction due to the loss of suitable habitats or changing climate conditions. By facilitating the movement of species to more suitable environments, it increases their chances of survival and reduces the risk of local extinctions.
- **Increased Biodiversity:** By introducing new species or expanding the range of existing species, it can increase species richness and promote more diverse ecosystems. This can contribute to ecosystem resilience and adaptive capacity.

POTENTIAL DISADVANTAGES:

- Species introductions may unknowingly introduce pests or diseases into new areas, particularly with longer transfer distances.
- Long-distance transfers based on projected climate conditions raise the likelihood that current habitat may not be suitable, which could result in poor growth or planting failure.
- Site factors other than climate, such as soil type, moisture regime, herbivory, competition, endemic pests and pathogens, and photoperiod may preclude successful establishment.
- Newly introduced species may become invasive.
- Can be resource-intensive and logistically challenging. It requires careful planning, monitoring, and long-term management to ensure the success of introduced populations. Limited resources and competing conservation priorities may make assisted migration impractical or less feasible in certain cases.
- Uncertainty in predicting future species' ranges and community compositions with climate disruption. The paleo-ecological record shows that current species assemblages break apart and new assemblages form when the climate changes.

ADAPTATION RESPONSE:

MANAGEMENT PRACTICES TO MAINTAIN / IMPROVE THE ABILITY OF FORESTS TO RESIST PESTS & PATHOGENS

CHARACTERISTICS:

Forest management practices that manipulate the density, structure, or species composition of a forest may reduce susceptibility to some pests and pathogens, inter alia:

- Thinning to reduce the density of a pest's host species to discourage infestation, based on the knowledge that species are especially susceptible to pests and pathogens at particular stocking levels.
- Adjusting rotation length to decrease the period of time that a stand is vulnerable to insect pests and pathogens, based on the knowledge that species are especially susceptible to pests and pathogens at particular ages.
- Creating a diverse mix of forest or community types, age classes, and stand structures to reduce the availability of host species for pests and pathogens.
- Managing canopy conditions depending on types of invasive species, for example, maintaining closed-canopy conditions to reduce the ability of light-loving invasive species to enter the understory or keeping canopy more open to reduce spreading of species (e.g. *Pinus strobus*) or pathogens that prefer conditions of shade, less wind, and higher humidity.
- Using biological control methods to manage pest populations in heavily infested areas.
- Restricting harvest and transportation of logs near stands already heavily infested with known pests or pathogens.
- Using impact models and monitoring data to anticipate the arrival of pests and pathogens and prioritize management actions.

PRIORITY TOPIC ADDRESSED:

5 INVASION BY NON-NATIVE SPECIES

MAIN RISKS ADDRESSED:

- Invasion by non-native species may result in biome shifts, with consequent changes in the spectrum of forest ecosystem services provided

INTENDED EFFECTS:

Improved non-native species management with dedicated measures for prevention, early detection, control management, including rapid response and rehabilitation and restoration.

KEY ADVANTAGES & POTENTIAL DISADVANTAGES:

Depends on management practices.



ADAPTATION RESPONSE:

CLOSE-TO-NATURE SILVICULTURE / ECOSYSTEM- BASED MANAGEMENT

CHARACTERISTICS:

Treating forests as an ecological system that performs multiple functions. This approach provides an array of ecosystem services with the aim of promoting forest health and biodiversity while also producing timber and other forest products. The approach is based on the principle that forests are complex ecosystems that function best when natural processes from the point of view of mass and energy fluxes are continuous and automatic.

Uses practices designed to create forest landscapes that contain all the diversity and irregularity of natural forests, for example with regard to the size and shape of stands and the presence of trees of different sizes. Forest management interventions are designed to mimic natural disturbances such as windthrow, insect outbreaks, and fire. This can include measures such as selective thinning, gap creation, and retention of deadwood and snags towards creating a more diverse forest structure.

Involving local communities in forest management can help ensure that forest management practices are adapted to local needs and priorities, while also promoting sustainable use of forest resources. This can include measures such as community-based forest management agreements, forest user groups, and forest certification schemes.

PRIORITY TOPIC ADDRESSED:

FOREST ECOSYSTEM SERVICES

MAIN RISKS ADDRESSED:

- Alteration of critical ecosystem services, including provisioning services (bioenergy, water); regulating services (climate regulation, carbon sequestration and storage, flood control, and pollination, pest and disease control); supporting services (primary production like timber); and cultural services (recreation, aesthetic and spiritual benefits)

INTENDED EFFECTS:

Improved forest health and biodiversity conservation.

Maintaining a diverse forest structure and reducing the risk of large-scale disturbances such as wildfires can help improve forest health, promote biodiversity, and enhance resilience to climate change.

Sustainable timber production.

Various continuous cover and selection harvesting practices, including close-to-nature silviculture, can provide a steady supply of timber while also maintaining forest health and biodiversity.

Carbon sequestration.

Maintaining healthy forests and forestry practices that emphasize development of high biomass forests can help sequester carbon and mitigate the effects of climate change.

Improved aesthetics and recreational value.

Creating a more natural forest structure, close-to-nature silviculture can improve the aesthetics of the forest and provide opportunities for recreational activities and sustainable tourism.



KEY ADVANTAGES:

- By imitating natural disturbance patterns, such as individual tree mortality or gap dynamics, it allows for the regeneration and development of diverse age classes and species composition. This promotes forest adaptability and improves long-term ecosystem stability.
- Through selection harvesting individual or small groups of trees, it can provide a continuous supply of high-quality timber while minimizing the negative impacts on forest ecosystems.
- By maintaining a diverse range of tree species and age classes, forests under this management approach can sequester and store more carbon after conversion to site-endemic species compositions and development of high biomass conditions. This contributes to climate change mitigation by reducing greenhouse gas emissions and enhancing carbon sinks.
- There is an opportunity to further expand the portfolio of close-to-nature silvicultural practices, for instance through greater emphasis on large dead tree retention and downed wood enhancement, irregularly structured and shaped gap creation, development of patch mosaics, and incorporation of tip-up mounds into managed stands.
- Close-to-nature silviculture can be used as a “conversion” or restoration method to transition plantations or monocultures to site-endemic species compositions, including species mixtures that are more adapted to future climatic conditions.

POTENTIAL DISADVANTAGES:

- Complexity and knowledge requirements: Forest managers need to have a good understanding of the natural dynamics of the forest, including tree species interactions, regeneration strategies, and long-term planning.
- Reduced timber production: Overall timber yield may be lower compared to intensive plantation forestry or clear-cutting practices. Longer rotation periods between harvests compared to traditional management systems may pose financial challenges for forest owners or managers who rely on timber income.
- Management costs associated with this approach can be higher compared to more conventional management systems. The need for continuous monitoring, selective harvesting, and maintenance of desired forest structures can increase the overall management expenses.
- Limited applicability: May not be suitable for all forest types or conditions. It is best suited for certain forest ecosystems with specific ecological characteristics. In some cases, the natural dynamics of the forest may not support continuous cover management, or the economic and social context may not favor this approach.
- Closer-to-nature forestry practices which promote the retention of deadwood in forests might, depending on the local conditions, require further precautions to prevent forest fires or undesired insect infestations.



ADAPTATION RESPONSE:

INTEGRATING FOREST-WATER INTERACTIONS INTO ADAPTIVE FOREST MANAGEMENT

CHARACTERISTICS:

Adaptive forest management options to manage hydrological impacts of climate change include:

- Thinning to promote drought resistance and improved forest growth.
- Removal or control of invasive species.
- Leaving a certain proportion of dead wood in forests, arranged in a slanting or perpendicular pattern relative to slope orientation.
- Removal of infected trees, and pest control by introducing predators and biocontrol agents.
- Increasing water retention facilities (e.g., terraces and ponds) to guarantee water supply and restore aquatic and floodplain habitats.
- Watershed-level planning to incorporate considerations of the hydrological processes and their interaction with the forest ecosystem. This approach will help identify critical areas that need protection and prioritize management actions accordingly.
- Maintaining drainage of forest roads, use a net of trenches for water retention via polders and keep undamaged soil and humus layer as a protection against erosion. Installing a greater frequency of water bars (shallow ditches placed diagonally across forest roads), larger culverts, and enhanced stream crossing structures.
- Reducing forest road densities and improving the design, layout, and engineering of the permanent forest road system. This approach may conflict with certain styles of close-to-nature forestry that rely on dense road or skid trail networks. Therefore, assessment of tradeoffs and multicriteria optimization of road network design is important.
- Promoting agroforestry practices, including the integration of trees into agricultural landscapes. Benefits include improved soil health, increased water infiltration, and reduced erosion. Promoting agroforestry practices in areas adjacent to forests can help reduce pressure on forests for agricultural land and promote sustainable land use practices.
- Maintaining riparian zones to act as buffers between the water body and forest. They help reduce erosion and sedimentation, maintain water quality, and provide habitat for aquatic species.

PRIORITY TOPIC ADDRESSED:

7 FOREST – WATER INTERACTIONS, INCLUDING HYDROLOGIC REGULATION AND RIPARIAN DYNAMICSS

MAIN RISKS ADDRESSED:

- Altered hydrology regimes due to climate change affecting water quality, aquatic habitats, and species and soil resources
 - Reduced water uptake by trees and reduced soil infiltration from large-scale disturbances, such as wildfire, bark beetle outbreaks and defoliating insects
 - Increased runoff and severe erosion and chemical loading
 - Accelerated rate of nutrient cycling, promoting increased forest growth and elevated nitrogen levels in streams
-
- Implementing silvopastoral systems that integrate tree cover into pasture lands. Silvopasture helps reduce soil erosion, improve soil fertility, and provide shade and shelter for livestock. These systems can also help increase water infiltration and recharge groundwater.

INTENDED EFFECTS:

Protected and improved water quality by minimizing soil erosion, reducing sedimentation in water bodies, and filtering pollutants.

Adaptive management practices such as riparian buffer zones, reforestation of degraded areas, improved forest road engineering and maintenance, and sustainable logging techniques can help maintain or restore water quality, stream habitats and increase flood resilience.

Increased water availability and flow regulation.

Preserving or restoring riparian vegetation for example can help stabilize streambanks, reduce water loss through evapotranspiration, and regulate water flow during periods of high rainfall or drought.

Watershed resilience.

Restoring and promoting healthy, well-functioning watersheds is important to maintain overall resilience in the face altered disturbance regimes and vegetation shifts.



KEY ADVANTAGES:

- Enhanced ecosystem services including wood production and improved water availability, reduced erosion, enhanced water purification, and increased flood regulation, benefiting both natural ecosystems and human communities.
- Supports the principles of integrated water resources management and promotes water security.

POTENTIAL DISADVANTAGES:

- Requires a comprehensive understanding of forest ecology, hydrology, and water resource management. It can be a complex undertaking, requiring expertise and coordination among different stakeholders, including forest managers, water authorities, and scientists. Adequate knowledge, data, and monitoring systems are essential for effective integration.
- Balancing multiple objectives within the forest-water nexus can present challenges and trade-offs. For example, maximizing timber production may conflict with maintaining water quality or protecting sensitive aquatic ecosystems. Integrating these objectives requires careful consideration and decision-making processes to ensure that all aspects are adequately addressed. Related planning and implementation challenges may arise due to limited resources, competing land uses, governance issues, and conflicting stakeholder interests.
- Forest and water resource dynamics are often slow to respond to management interventions, and the benefits may take time to materialize fully.





5 Relevant Initiatives and Actors

5.1 INTRODUCTION

This chapter outlines ongoing initiatives that complement the aforementioned climate adaptation pathways. These initiatives include notable projects, published databases and remote sensing platforms, research programs, and protective forest policies. Together, these efforts aim to enhance forest resilience and adaptive capacity in the face of climate change.

5.2 RELEVANT PROJECTS

CARPATCLIM – CLIMATE OF THE CARPATHIAN REGION

[CARPATCLIM \(2010-2013\)](#) brought together 10 research institutes from Austria, Croatia, Czechia, Hungary, Poland, Romania, Serbia, and Ukraine with funding from the European Commission to “improve the basis of climate data in the Carpathian Region for applied regional climatological studies such as a Climate Atlas and/or drought monitoring, to investigate the fine temporal and spatial structure of the climate in the Carpathian Mountains and the Carpathian basin with unified methods.” Assessing the

1961 to 2010 timeframe, its three work modules produced a comprehensive understanding and homogenization of climate data for the larger Carpathian region, publishing these as online Atlases, gridded climatologies, publicly available datasets, and drought monitoring. As such, the resources CARPATCLIM made available online remain a vital resource for climate impact assessment, forecasting, and planning. The project remains a model for region-wide monitoring based on data sharing and standardization.

CLIMATE SMART FOREST MANAGEMENT FOR CENTRAL AND EASTERN EUROPE

A promising seven-year project (2020 – 2027), called [Climate-Smart Forest Management for Central and Eastern Europe \(CLIMAFORCEELIFE\)](#), is developing, demonstrating, and promoting climate-smart management techniques for Carpathian forests and beyond. Starting with demonstration sites in 3 countries (Bulgaria, Hungary, and Slovakia), the aim of this €5.6 million project is to encourage the transition to future-adaptive forestry across the Central and Eastern European region. Based on extensive stakeholder involvement, the project will transfer experience and information to the forest sector through guidance documents and a variety of science communication media. A strength of the project is its grounding in “real-world” adaptive (or climate-smart) forestry operations, meaning the demonstration sites are located on managed forests where techniques are applied and tested actively. This will allow information to be developed that is readily transferrable and relevant to diverse stakeholders. Techniques such as close-to-nature silviculture, retention forestry, control of herbivore browsing, large woody debris enhancement, and gap-based harvesting are adapted to local conditions, anticipating future climate-driven shifts in forest composition and productivity as well as disturbance risks.

The CLIMAFORCEELIFE project was launched at an opening conference held in Bratislava, Slovakia in October of 2021, organized as part of the EU Strategy for the Danube Region 10th Annual Forum. The event brought together over 130 forest experts, scientists, and stakeholders from across the greater region, resulting in a large number of presentations and resources about climate change and forest adaptation

in the Carpathian region [that are now available online](#). CLIMAFORCEELIFE recently partnered with a project funded by the Trust for Mutual Understanding (W.S. Keeton and R. Aszalos P.I.s) called Bridging the Divide between European and North American Perspectives on Ecological Silviculture. The latter initiative has conducted several scientific exchange events in Romania, Hungary, Slovakia, Czechia, and the United States of America. These events facilitated the sharing of experiences and information around climate adaptive forest management from the Carpathian and U.S. perspectives.



Workshop for forestry practitioners on close-to-nature silviculture and climate-smart forest management, held in the Börzsöny mountains of Hungary, June 2023. The event was organized by the Centre for Ecological Research, WWF Hungary, and the University of Vermont with funding from CLIMAFORCEELIFE and the Trust for Mutual Understanding.

5.3 RELEVANT TOOLS

PAN-EUROPEAN FOREST RISK KNOWLEDGE MECHANISM

Building on the [Bratislava Ministerial Resolution “Adapting pan-European forests to climate change” \(2021\)](#), FOREST EUROPE is working towards the pan-European forest risk knowledge mechanism (FoRISK) to support adaptation of forests to changing climatic and site conditions and enhance forest resilience at a pan-European level. The vision of the FoRISK is to provide relevant evidence-based forest risk and

adaptation-related information to political decision-makers based on trustful cooperation with scientists, practitioners, and society.

The FoRISK Pilot, running from September 2022 to February 2024, tests policy tools for various disturbance factors like wildfires, biotic threats, and storms. The Pilot involves National Focal Points from the Carpathian Convention dealing

with forestry issues to evaluate the feasibility, funding options, adjustments, and improvements related to the FoRISK concept. The Pilot has three successive but interacting pilot phases, each with a specific thematic focus on a forest damage agent and a comparable approach [Pilot phase 1: Wildfires (9/2022-3/2023); Pilot phase 2: Pests & diseases (3/2022 – 8/2023); Pilot phase 3: Storms (9/2023 – 2/2024)].

EUROPEAN PRIMARY AND OLD-GROWTH FOREST DATABASES

The current EU Biodiversity Strategy sets an ambitious goal of conserving all remaining primary and old-growth forests by the year 2030. Achieving this goal relies on the availability of accurate old and natural forest inventory (i.e. spatial) data for the entire EU, which presently includes all Carpathian countries except Ukraine and Serbia (both have EU candidate status). Several completed and ongoing assessments have addressed this need. In these initiatives, the term “primary forest” is based on the FAO definition (FAO 2018), whereas old-growth is defined differently by the European Commission (2015), [as described in the new EU guidelines adopted by the European Commission in 2023](#). The datasets produced by these assessments will be vital for informing conservation planning, structurally complex forest restoration, adaptive management, and regional forest carbon management strategies, as primary and old-growth forests in the Carpathians typically have exceptionally high carbon stocking (Keeton et al. 2010; Burrascano et al. 2013).

Two assessments of particular relevance are those by Sabatini et al. (2018, 2020) and Barredo et al.

(2021). The former assembled a comprehensive data base spanning all previously documented and mapped primary forest sites across 32 countries. A predictive model was then developed from this dataset to estimate the likely occurrence and distribution of additional, unmapped primary forests. The pan-European database was later expanded to incorporate 51 unique data sets from 35 countries, importantly including Russia (Sabatini et al. 2021). This increased the estimated area of primary forest from 1.4 to 41.2 million ha, a massive expansion due largely to the inclusion of Russia and Norway. The state of known, ground validated primary and old-growth forest occurrences was further summarized and updated in Barredo et al. (2021).

The Carpathian Convention has been establishing an [Inventory of Virgin Forests of the Carpathians](#), based on officially reported data by the Parties, which shall strengthen the protection of one of the last remaining virgin forests in Europe. This has been undertaken in cooperation with the European Environment Agency (EEA), the European Topic Centre on Urban, Land, and Soil Systems (ETC-ULS).

The current inventory was welcomed by the Carpathian Convention COP7 in October 2023. In addition, this inventory was extended to other degrees of naturalness, particularly quasi-virgin forests in the Carpathians, and specific criteria and indicators for the selection of quasi-virgin forests in the Carpathians were finalized and adopted at COP7. These data come from national inventories and often produce different estimates as compared to other sources (see, for example, Mikolas et al. 2019). All of the aforementioned datasets include the most up-to-date spatial data for primary and old-growth forests in the Carpathians, although downscaling, accuracy assessment, and correction of inventory gaps are ongoing.



DISTURBANCE MONITORING AND RESPONSE CAPABILITY

Disturbance risk monitoring and prediction will be critical elements of adaptive forest management in the Carpathians. This point was clearly stressed by the regional experts surveyed in this assessment and is at the forefront of on-going scientific research.

There are promising technological advances that will greatly aid this endeavour. These include several remote sensing platforms used to track and monitor disturbance events and disturbance risks, often in real time depending on the platform.

For example, the [Global Ecosystem Dynamics Investigation \(GEDI\)](#) provides high resolution, three-dimensional laser ranging of the Earth's forests, enabling assessment of canopy structure and fuel profile that can be used to map fire risks. The satellite mounted [MODIS \(Moderate Resolution Imaging Spectroradiometer\)](#) detects day and nighttime thermal anomalies on the Earth's surface. This allows users to monitor fire occurrences day to day for any region of the world.

As another example, the EU has invested in the Structured Approaches for Forest Fire Emergencies in Resilient Societies (SAFERS). It combines remotely sensed data from the Copernicus and the Global Earth Observation System of Systems (GEOSS) platforms with crowd sourced data to create "[an open and integrated platform featuring a forest fire decision support system.](#)" The wildFire cLimate impacts and Adaptation Model (FLAM), hosted by IIASA, operates with a daily time step at various spatial resolutions and uses mechanistic fire modeling algorithms to parameterize the impacts of climate, fuel availability, and human activities on wildfire probabilities, frequencies, and burned areas. FLAM uses daily climate data for temperature, precipitation, wind, and relative humidity. Its modular



structure allows for the inclusion of additional fire-related variables such as distance to roads, cropland, lakes, slope, and elevation. The model is therefore adaptable to unique regional characteristics such as traditional practices of agricultural burning, forest clearing, or peatland draining. Eventually this will allow for the production of integrated regional hot-spot maps for wildland fire risk. One case study implemented with FLAM focussed on current wildland fire patterns and challenges in Europe (Fernandez-Anez, N. et al. 2021).

Remote sensing platforms are providing valuable insights into the changing environmental conditions across the Carpathian region and Europe generally. An example is the use of Sentinel-2 imagery to derive changes in tree phenology (timing of spring bud break) in Poland (Grabska-Szwagrzyk & Tymńska-Czabańska 2023). This method has proven effective for monitoring interspecific variation in phenological responses to climate change, with implications for forest productivity, carbon budgets, hydrologic regulation, and energy fluxes. Other technologies are available to assist forest managers in predicting disturbance risks into the future under different climate and forest management scenarios. Examples include a variety of sophisticated models that simulate disturbances and their interactions

with forest dynamics for millions of trees or thousands of pixels across entire landscapes (see, for example, Seidl et al. 2012; Kruhlov et al. 2018). Combining these technologies will give institutions, scientists, and forestry practitioners advanced capabilities, allowing them to better anticipate, plan, and respond to disturbances. Perhaps even more importantly, disturbance prediction capabilities will help forest managers strategically plan the type and spatial configuration of forest restoration and adaptive silvicultural interventions used to reduce or limit disturbance risks, such as efforts to increase forest landscape heterogeneity.

As important as disturbance risk prediction is enhancing the capacity to manage and respond to emerging threats, such as the growing forest fire risk in some parts of the Carpathian region. Carpathian counties are expanding these capabilities through a variety of funding mechanisms and cooperative programs. These include the EU's initiatives to assist Member States in forest fire prevention (rather than suppression) and cooperative programs between the United States Forest Service and variety of a ministries and stakeholders in Ukraine to improve forest fire management capacity through training and acquisition of fire-fighting resources.

RESEARCH

5.4 RELEVANT RESEARCH INITIATIVES

EUROPEAN WIDE FOREST DISTURBANCE MAPPING AND PROJECTION MODELING

The [Ecosystem Dynamics and Forest Management group](#) at the Technical University of Munich, Germany, led by Dr. Rupert Seidl, is conducting ongoing research on climate change effects on forests across Europe. This includes the use of remotely sensed data to map and track changing disturbance regimes and the interactions among disturbance agents. Related studies using simulation models project disturbances and their consequences for specific landscapes and the European continent under alternate

climate scenarios. [This group has published numerous papers](#) on disturbed dynamics, climate impacts on biodiversity, effects on ecosystem services, and forest resilience that are directly relevant to the Carpathian region. This body of research now comprises an indispensable resource for understanding and predicting the consequences of climate change for European forests as well as opportunities for adaptive forest management.



SUPERB - UPSCALING FOREST RESTORATION / EUROPEAN FOREST INSTITUTE

Disturbance risk monitoring and prediction will be critical elements of The European Forest Institute (EFI), an international organization with offices in 41 countries, has taken a lead role in producing sustainable forest management guidance that is directly relevant to several of the key risks and adaptation responses identified in this assessment. Working closely with research organizations, forestland owners, and industry, EFI produces information that is practical and timely in terms of addressing hot button forest management concerns.

Of particular relevance to climate risks and adaptation is EFI's Resilience Programme, which has produced a wide variety of reports offering guidance on climate data for European forestry, old-growth forest conservation, adaptive management in response to bark beetle outbreaks, disturbance trends and vulnerabilities, and state-of-the-art thinking on "closer" to nature silviculture. [Reports and information are made publicly available on the EFI website.](#)

EFI's Resilience Programme launched a large-scale project called SUPERB - Systemic solutions for upscaling of urgent ecosystem restoration for forest-related biodiversity and ecosystem services in partnership with Wageningen Environmental Research. Funded by a €20 million grant from the EU Horizon 2020 Research and Innovation Programme, combined with ["in-kind contributions of €90 million from its associated partners,"](#) SUPERB will synthesize state-of-the-art knowledge and examples (both successful and unsuccessful) of forest ecosystem restoration and adaptation relevant to a variety of contexts across Europe. At the core of SUPERB is a demonstration network consisting of 12 sites in 12 countries, one of which is located in the Făgăraș Range of the Carpathian Mountains in Romania. The Romanian demonstration site is testing restoration forestry practices for phased-conversion of spruce monocultures to mixed-species, structurally complex stands in buffer areas around core zones protecting old-growth forests. [A complete description is available online.](#)

POLICIES

5.5 RELEVANT POLICIES

PROTECTION OF VIRGIN FORESTS IN UKRAINE AND SLOVAKIA

In 2017 and in accordance with the Carpathian Convention, the Ukrainian government set an example for the region by establishing new legal protections for forest natural monuments. The law effectively called for the protection of all remaining virgin or natural forests. [According to WWF - Ukraine](#), 37 forest natural monuments had been created by 2019, totalling 4,750.8 ha of protected virgin and natural forests in the Trans-Carpathian and Ivano-Frankivsk oblasts (provinces). Since then, the area of newly protected natural, quasi-virgin, and virgin forest has expanded even further to more than 25,000 hectares.

The Slovak Republic followed suit not long afterwards. In November 2021, [the Slovak government approved establishment of a network of protected natural and virgin forest areas](#) comprising 6,500 ha distributed across 76 sites. The new reserve network, called Old Growth Forests of Slovakia, came in response to recently completed mapping

(Mikolas et al. 2019) showing that [“approximately 10,180 ha of old growth forests remained in Slovakia, of which one third was unprotected or insufficiently protected.”](#)

The protections in Ukraine and Slovakia both followed the establishment of a UNESCO World Heritage transnational network in 2007 entitled, “Ancient and Primeval Beech Forests of the Carpathians and Other Regions of Europe.” [This network now encompasses](#) 94 separate forested tracts in 18 countries, including 70 tracts in countries signatory to the Carpathian Convention. These protected forests are places where natural processes will predominate and source populations will be maintained, providing the core reserve component a holistic and multi-pronged forest adaptation approach. Furthermore, the protections recognize the necessity of safeguarding rare and unique natural communities and biodiversity hotspots to ensure their future persistence.

EU FOREST STRATEGY FOR 2030

The new EU Forest Strategy for 2030 (EC 2021) is one of the flagship initiatives of the European Green Deal and builds on the EU Biodiversity Strategy for 2030. It recognises the central and multifunctional role of forests, and the contribution of foresters and the entire forest-based value chain for achieving a sustainable and climate neutral economy by 2050 and preserving lively and prosperous rural areas. The EU Forestry Strategy sets a vision and proposes concrete actions to improve the quantity and quality of EU forests and strengthen their protection, restoration and resilience. It aims to adapt Europe’s forests to the new conditions, weather extremes and high uncertainty brought about by climate change. This is seen as a precondition for forests to continue delivering their socio-economic functions. Provisions of the strategy cover aspects of protecting, restoring and enlarging the EU’s forests to combat climate change, reverse biodiversity loss and ensure resilient and multifunctional forest ecosystems. This includes, inter alia, promoting a sustainable forest bioeconomy, protecting EU’s last remaining primary and old-growth forests, and ensuring forest restoration and reinforced sustainable forest management for climate adaptation and forest resilience.


In March 2023 the European Commission adopted two sets of guidelines to support the implementation of the forest strategy within the broader framework of the EU Biodiversity Strategy for 2030, while supporting the

general EU commitments under the Kunming-Montreal Global Biodiversity Framework (Convention on Biological Diversity). These guidelines shall support efforts to strengthen the protection, restoration and resilience of EU forests to help them adapt to a changing climate and improve their quantity and quality. The Guidelines on Biodiversity-Friendly Afforestation, Reforestation and Tree Planting provide a set of practical recommendations to support authorities, forest and landowners, and managers and civil society to better implement biodiversity-friendly afforestation, reforestation and tree-planting projects including at the local level. Through both active planting and natural regeneration, these guidelines constitute one of the key milestones to implement the 3 billion additional trees pledge of the EU by 2030. They address afforestation initiatives in agricultural land; reforestation actions in forest land, including restoration actions; and tree planting in urban and peri-urban environments, as well as agricultural land (agroforestry). The Guidelines for Defining, Mapping, Monitoring and Strictly Protecting EU Primary and Old-Growth Forests provide practical guidance to national policy- and decision-makers that will allow them to effectively identify and protect remaining primary and old-growth forests in the EU. The guidelines set out criteria for identifying primary and old-growth forest areas based on a list of indicators or principles. A timeline is suggested for their mapping and strict protection.



6

Conclusions and Recommendations



6.1 INTRODUCTION

The Carpathian Mountains are a region of significant ecological and cultural importance, which is under threat from climate change and anthropogenic disturbances. The impacts of climate change on forests in the Carpathian Mountains are expected to lead to changes in forest composition, structure, and function, as well as potential feedbacks to the climate system.

To address these impacts, a range of adaptation and mitigation strategies have been proposed, which aim to reduce the vulnerability of forest ecosystems to climate change and reduce greenhouse gas emissions from forest management activities. The implementation of these strategies will require cooperation and coordination among stakeholders, including forest managers, policymakers, and local communities.

The vulnerability of Carpathian forests to climate change poses significant challenges to the ecological and socio-economic functions, as well as to the livelihoods of forest-dependent communities.

However, through adaptive forest management practices, cross-boundary collaboration, stakeholder engagement, and the integration of local knowledge, Carpathian forests can become more resilient and better equipped to cope with the impacts of climate change. Addressing research gaps and improving long-term monitoring and data availability can also help inform evidence-based adaptation strategies, while ensuring that policies and frameworks explicitly consider climate change impacts in long-term planning and regional coordination.

The following highlights opportunities and pathways in this regard as well as further research needs for evidence-based decision-making.

6.2 OPPORTUNITIES AND PATHWAYS

Given the unique ecological and socioeconomic characteristics of the Carpathians, there are several key pathways to further consider for climate-resilient forest management practices. Fostering collaboration and transboundary cooperation among the Carpathian countries will be vital for effective climate change adaptation in forestry, especially through sharing knowledge and experiences with various approaches and developing pathways for addressing common challenges.

Considering the history of deforestation in the Carpathians, one key pathway is to focus on forest restoration and reforestation efforts. This involves restoring degraded forest areas, establishing new close-to-nature forests, promoting natural regeneration of forests, converting planted monocultures to the site-endemic and/or future-adapted species compositions, and protecting and reintroducing rare native tree species in their natural ecosystem, following single or a combination of adaptation response options outlined in Chapter 5.

Protect and Conserve Natural Forests:

This is crucial for climate change adaptation in the Carpathians. Establishing and effectively managing protected areas, national parks, and nature reserves helps preserve intact forest ecosystems, maintain biodiversity, and provide refuges for species. These protected areas also contribute to carbon sequestration and storage.

Enhance Forest Landscape Connectivity:

This is vital for allowing species to migrate and adapt to changing climate conditions. This can be achieved through the creation of ecological corridors, which connect fragmented forest patches, enabling species movement and facilitating gene flow. Conserving and restoring riparian zones can also enhance connectivity and provide climate refugia for species.

Manage for Diverse Landscape Composition and Structure:

This will help make forest ecosystems more resilient to disturbances, particularly bark beetles and other insect pests as well as fungal pathogens. Increasing the heterogeneity of forest patches both within stands and across landscapes limits contagion and dampens the ability of insects to disperse and colonize new host trees.

Adopt Ecosystem-based Adaptation:

This involves recognizing and harnessing the benefits provided by intact ecosystems and promoting sustainable livelihoods in an inclusive approach. This includes conserving and restoring wetlands, protecting carbon-rich forest ecosystems, coordinated invasive species control, and integrating nature-based solutions such as green infrastructure and natural water retention measures as blue infrastructure.

Forest Fire Management, Prevention, and Restoration:

As climate change increases the risk of high intensity forest fires in the Carpathians, implementing effective forest fire management and prevention measures is critical. This pathway involves developing national and regional early warning systems, improving fire suppression capabilities, and promoting community-based fire management approaches. Ensuring adequate resources and training for forest fire management is essential to mitigate the impacts of wildfires on forests and communities. In forest types where low intensity fires were historically an important part of the natural disturbance regime, silvicultural techniques like fuels treatment and prescribed burning will be useful to restore stand structures that will once again support beneficial ground fires, while limiting the spread of crown fires.

“Fostering collaboration and transboundary cooperation among the Carpathian countries will be vital for effective climate change adaptation in forestry, especially through sharing knowledge and experiences...”

Promote Sustainable Wood Utilization and Develop Value Chains for Forest Products:

These efforts can enhance the economic viability of forests while supporting climate change adaptation. This pathway involves encouraging responsible harvesting practices, supporting local processing industries, and promoting the use of sustainably sourced wood products.

Sustainable Forest Practices and Certification:

Adopting sustainable forestry practices, such as continuous coverforestry and reduced impact logging techniques, help minimize ecosystem disturbance and maintain forest health. Certification for sustainable forest management may improve market access and support livelihoods. Experience has been mixed as to whether certified wood products are able to generate price premiums.

Education and Capacity Building:

Building the capacity of forest managers, policymakers, local communities, and relevant stakeholders is a key pathway for climate change adaptation. This includes providing training, education, and awareness campaigns to enhance understanding of climate change impacts, adaptation strategies, and the importance of sustainable forest management.

Harmonize Monitoring:

Monitoring climate forecasts, disturbance risks, and biodiversity threats will help the region coordinate a variety of adaptation measures, both conservation oriented and responsive to new threats as they arise.

Incorporate Traditional and Local Knowledge:

This can provide valuable insights into forest dynamics, species behavior, and ecosystem responses to climate change. Community involvement also promotes local ownership, improves livelihoods, and fosters sustainable forest practices.

6.3 KNOWLEDGE GAPS AND RESEARCH NEEDS

Closing the following knowledge gaps will improve our understanding of climate dynamics and adaptation in forestry and enable the development of effective and context-specific strategies to enhance the resilience of forest ecosystems to climate change. Collaboration between researchers, practitioners, landowners, and stakeholders is crucial to address these gaps and ensure that adaptation efforts are science-based, practical, and inclusive.

IMPROVED REGIONAL-SCALE FOREST MONITORING:

Forest monitoring is currently primarily under the purview of ministries and research institutions within individual Carpathian countries. Consequently, the region lacks a coordinated monitoring network, important for forecasting and responding to vulnerabilities, such as forest fire risks, as they develop. One such initiative was the CARPATCLIM project, a joint initiative to develop standardized climatological databases for the Carpathian region. Concluded in 2010 (see Chapter 5), the initiative provided a model that could be extended to other areas of importance within the forest sector, such as disturbance risks, biodiversity, and forest health. Critical questions remain to be resolved, such as the priorities for regional monitoring, their relative value, and their scope. Clear benefits to the region regarding climate forecasting, planning, conservation, and adaptation would need to be established.


If monitoring were further coordinated to enable standardization of monitoring parameters and data at the regional level, it would facilitate both joint adaptation efforts, such as multilateral plans for assisted species migration, and enable comparison of research results across the Carpathian ecoregion (Grodzki et al. 2018). Regional forest monitoring could be expanded to include additional layer and indicators for forest ecosystem dynamics under climate change, including shifts in forest disturbances, structure, species distribution patterns, and ecosystem functioning. Furthermore, forest ecosystems can exhibit ecological thresholds, beyond which they may experience rapid and irreversible changes in structure and function. However, our understanding of these thresholds and the conditions that trigger them is limited and should be further investigated and monitored.

IMPORTANCE OF GENETIC DIVERSITY:

Another area for additional knowledge generation concerns the importance of genetic diversity in forest ecosystems for adaptation. This involves studying the genetic characteristics of tree species, assessing the adaptive potential of different genetic lineages, and investigating how genetic diversity influences ecosystem resilience and productivity.

ADAPTIVE CAPACITIES OF FORESTS:

There is also a need for more comprehensive research on how different tree species and forest ecosystems respond to changing climatic and soil conditions. This includes understanding the physiological and phenological responses of trees and the interactions between species and their environment. While some research has been conducted on the adaptive capacity of forests, there are still knowledge gaps in understanding the mechanisms by which forests can adjust their composition, structure, and functioning to changing conditions, as well as the limits to their adaptive capacity.




EFFECTIVENESS OF SILVICULTURE PRACTICES:

Further research would also be needed for assessing the effectiveness of various adaptive silviculture practices in Carpathian forests. This includes evaluating the effects of different adaptation approaches (such as those outlined in Chapter 4) on forest resilience, productivity, biodiversity and socio-economic factors. Long-term monitoring of adaptive practices will also be important to continuously (re-)evaluate their success. Research is further needed to investigate the interactions and synergies among different adaptation options. Understanding how different approaches can complement or conflict with each other is crucial for optimizing adaptation strategies and avoiding unintended consequences.

SOCIOECONOMIC EFFECTS AND IMPACTS:

Another critical knowledge gap relates to understanding the socio-economic dimensions of adaptation in forestry. This includes assessing the economic viability and costs associated with different adaptation approaches, understanding the social acceptability and equity implications of adaptation interventions, and considering the impacts of adaptation on local communities and livelihoods. On the latter, more research and case studies are needed to foster the integration of traditional ecological knowledge into adaptation strategies, recognizing the importance of local communities' perspectives, practices, and governance systems in sustainable forest management.



6.4 CONCLUSION

This assessment relied primarily on a qualitative survey to gauge concerns about climate change and Carpathian forests expressed by relevant ministry-based and academic professionals. As such, the report's conclusions are best used to complement other types of studies and assessments, such as those using quantitative projections and spatial analyses of climate risks and vulnerabilities.

And yet, the report's findings show clear commonality among experts in the Carpathian countries in terms of key areas of concern. These transcend long-standing concerns like climate change effects on forest productivity and species composition. Rather, there are emerging and serious questions around risks posed by drought, floods, insects, forest fires, invasive species, and other symptoms of global change now being experienced strongly throughout the Carpathian mountain region. In many ways, these risks are bringing to light new vulnerabilities in the forest sector, making adaptive approaches for both sustainable forest management and biodiversity conservation more urgent than ever. The key messages reported here—about both risk reduction and adaptive management—will help shape continued dialogue around multi-lateral planning and cooperation in the face of climate change.

REFERENCES

- Alberton, M., Andresen, M., Citadino, F., Egerer, H., Fritsch, U., Götsch, H., Hoffmann, C., Klemm, J., Mitrofanenko, A., Musco, E., Noellenburg, N., Pettita, M., Renner, K., Zebisch, M. (2017). Outlook on climate change adaptation in the Carpathian mountains. United Nations Environment Programme, GRID-Arendal and Eurac Research. Nairobi, Vienna, Arendal and Bolzano. www.unep.org, www.grida.no, www.eurac.edu
- Aszalós, R., Thom, D., Aakala, T., Angelstam, P., Brümelis, G., Gálhidy, L., Gratzer, G., Hlásny, T., Katzensteiner, K., Kovács, B., Knoke, T., Larrieu, L., Motta, R., Müller, J., Péter Ódor, Rozenberger, D., Paillet, Y., Pitar, D., Standovár, T., Svoboda, M., Szwagrzyk, J., Toscani, P., Keeton, W.S. (2022). Natural disturbance regimes as a guide for sustainable forest management in Europe. *Ecological Applications* 32: e2596
- Baradat, E., Hlásny, T., & Merganičová, K. (2019). Forest adaptation to climate change in the Carpathian Mountains: A review. *Annals of Forest Science*, 76(2): 1-13.
- Barredo, J.I., Brailescu, C., Teller, A., Sabatini, F.M., Mauri, A., Janouskova, K. (2021). Mapping and assessment of primary and old-growth forests in Europe. EUR 30661 EN, Publications Office of the European Union, Luxembourg. ISBN 978-92-76-34230-4, doi:10.2760/797591, JRC124671.
- Bokwa, A., Klimek, M., Krzaklewski, P., Kukułka, W. (2021). Drought Trends in the Polish Carpathian Mts. in the Years 1991–2020. *Atmosphere* 12: 1259. doi.org/10.3390/atmos12101259
- Burrascano, S., Keeton, W.S., Sabatini, F.M., Blasi, C. (2013). Commonality and variability in the structural attributes of moist temperate old-growth forests: A global review. *Forest Ecology and Management* 291:458–479.
- Carpathian Convention (2014). Criteria and Indicators for identification of virgin forests in the Carpathians. http://www.carpathianconvention.org/tl_files/carpathiancon/Downloads/03%20Meetings%20and%20Events/COP/2014_COP4_Mikulov/Follow%20Up/DOC13_Criteria_Indicators_virginforests_FINAL_26SEP.pdf. Accessed 29 August 2023.
- CARPATCLIM (2010-2013). Online resource accessed 29 August, 2023. <http://www.carpatclim-eu.org/pages/about/>
- Ceballos, L. S., Kyriazopoulos, A. P., & Dieterich, M. (2018). Climate change adaptation strategies for forests and forest management in the Carpathian Mountains. *Forest Policy and Economics* 90: 1-11.
- Costa, H., de Rigo, D., Libertà, G., Houston Durrant, T., & San-Miguel-Ayanz, J. (2020). European wildfire danger and vulnerability in a changing climate: towards integrating risk dimensions, EUR 30116 EN.
- Didovets, J., Krysanova, V., Bürger, G., Snizhko, S., Balabukh, V., Bronstert, A. (2019). Climate change impact on regional floods in the Carpathian region. *Journal of Hydrology: Regional Studies* 22: 100590.
- Dobrowolska, D., Bytnerowicz, A., Omasa, K., Paoletti, E., Wieser, G., Mikuška, P. (2017). Vulnerability of forests in the Carpathian Mountains: A synthesis. *Environmental Pollution*, 231, 1241-1252.
- Drever, C.R., Cook-Patton, S.C., Akhter, F., Badiou, P.H., Chmura, G.L., Davidson, S.J., Desjardins, R.L., Dyk, A., Fargione, J.E., Fellows, M., Filewod, B., Hessing-Lewis, M., Jayasundara, S., Keeton, W.S., Kroeger, T., Lark, T.J., Le, E., Leavitt, S.M., LeClerc, M.E., Lemprière, T.C., Metsaranta, J., McConkey, B., Neilson, E., St-Laurent, G.P., Puric-Mladenovic, D., Rodrigue, S., Soolanayakanahally, R.Y., Spawn, S.A., Strack, M., Smyth, C., Thevathasan, N., Voicu, M., Williams, C.A., Woodbury, P.B., Worth, D.E., Xu, Z., Yeo, S., Kurz, W.A. (2021). Natural Climate Solutions for Canada. *Science Advances* 7 (23): eabd6034
- European Commission. (2015). Natura 2000 and Forests - Part II. Office for Official Publications of the European Communities, Luxembourg. Technical Report - 2015 - 088. p. 108.
- European Commission. COMMUNICATION FROM THE COMMISSION TO THE EUROPEAN PARLIAMENT, THE COUNCIL, THE EUROPEAN ECONOMIC AND SOCIAL COMMITTEE AND THE COMMITTEE OF THE REGIONS (2021). New EU Forest Strategy for 2030. COM(2021) 572 final. Brussels, Belgium. https://commission.europa.eu/document/download/cf3294e1-8358-4c93-8de4-3e1503b95201_en.
- FAO (2018). Terms and Definitions FRA 2020, Rome. p. 26.
- Fernandez-Anez, N., Krasovskiy, A., Müller, M., Vacik, H., Baetens, J., Hukić, E., Kapovic Solomun, M., Atanassova, I., Glushkova, M., Bogunović, I., Fajković, H., Djuma, H., Boustras, G., Adámek, M., Devetter, M., Hrabalíkova, M., Huska, D., Martínez Barroso, P., Vaverková, M.D., Zúmr, D., Jögiste, K., Metslaid, M., Koster, K., Köster, E., Pumpanen, J., Ribeiro-Kumara, C., Di Prima, S., Pastor, A., Rumpel, C., Seeger, M., Daliakopoulos, I., Daskalakou, E., Koutroulis, A., Papadopoulou, M.P., Stampoulidis, K., Xanthopoulos, G., Aszalós, R., Balázs, D., Kertész, M., Valkó, O., Finger, D.C., Thorsteinsson, T., Till, J., Bajocco, S., Gelsomino, A., Amodio, A.M., Novara, A., Salvati, L., Telesca, L., Ursino, N., Jansons, A., Kitenberga, M., Stivrins, N., Brazaitis, G., Marozas, V., Cojocar, O., Gumeniuc, I., Sfecla, V., Imeson, A., Veraverbeke, S., Mikalsen, R.F., Koda, E., Osinski, P., Castro, A.C. M., Nunes, J.P., Oom, D., Vieira, D., Rusu, T., Bojović, S., Djordjevic, D., Popovic, Z., Protic, M., Sakan, S., Glasa, J., Kacikova, D., Lichner, L., Majlingova, A., Vido, J., Ferk, M., Tičar, J., Zorn, M., Zupanc, V., Hinojosa, M., Knicker, H., Lucas-Borja, M.E., Pausas, J., Prat-Guitart, N., Ubeda, X., Vilar, L., Destouni, G., Ghajarnia, N., Kalantari, Z., Seifollahi-Aghmiuni, S., Dindaroglu, T., Yakupoglu, T., Smith, T., Doerr, S., Cerda, A. (2021). CURRENT WILDLAND FIRE PATTERNS AND CHALLENGES IN EUROPE: A SYNTHESIS OF NATIONAL PERSPECTIVES. *Air, Soil and Water Research* 14 e117862212110281. <https://doi.org/10.1177/11786221211028185>.
- Fleischer, P., Jr., Holko, L., Celer, S., Čekovská, L., Rozkošný, J., Škoda, P., Olejár, L., Fleischer, P. (2020). Carbon balance and streamflow at a small catchment scale 10 years after the severe natural disturbance in the Tatra Mts, Slovakia. *Water* 12: 2917. <https://doi.org/10.3390/w12102917>.
- FOREST EUROPE. (2015). Relevant terms and definitions used for the updated pan-European indicators for sustainable forest management. p. 33.
- Forzieri G., Girardello M., Ceccherini G., Mauri A., Spinoni J., Beck P., Feyen L., Cescatti A. (2020). Vulnerability of European forests to natural disturbances, EUR 29992 EN, Publications Office of the European Union, Luxembourg, ISBN 978-92-76-13884-6, doi:10.2760/736558, JRC118512

- Forzieri, G., Marco, G., Ceccherini, G. et al. (2021). Emergent vulnerability to climate-driven disturbances in European forests. *Nature Communications* 12: 108. <https://doi.org/10.1038/s41467-021-21399-7>.
- Füssel, H.M., Lourenço, T.C., Downing, C., Hildén, M., Leitner, M., Marx, A., & Prutsch, A. (2018). National climate change vulnerability and risk assessments in Europe, 2018. European Environment Agency. Copenhagen, Denmark.
- García-Duro, J., Ciceu, A., Chivulescu, S., Badea, O., Tanase, M.A., Aponte, C. (2021). Shifts in forest species composition and abundance under climate change scenarios in southern Carpathian Romanian temperate forests. *Forests*. 12(11):1434. <https://doi.org/10.3390/f12111434>
- Grabska-Szwagrzyk, E. & Tymińska-Czabańska, L. 2023. Sentinel-2 time series: a promising tool in monitoring temperate species spring phenology. *Forestry: An International Journal of Forest Research* cpad039, <https://doi.org/10.1093/forestry/cpad039>
- Grodzki, W., Nowakowska, J. A. & Jagodziński, A. M. (2018). Climate change impacts and adaptation strategies in forests of the Carpathian Mountains, Poland. *Forests* 9(7): 1-20.
- Hlásny, T., Merganičová, K., Bucha, T., & Konôpka, B. (2017). Climate change impacts and adaptation scenarios in forests: A case study in the Carpathian Mountains. *Journal of Environmental Management* 197: 70-83.
- Hlásny, T., Krokene, P., Liebhold, A., Montagné-Huck, C., Müller, J., Qin, H., Raffa, K., Schelhaas, M.-J., Seidl, R., Svoboda, M., Viiri, H. (2019). Living with bark beetles: impacts, outlook and management options. *From Science to Policy* 8. European Forest Institute.
- IPCC. (2014). *Climate Change 2014: Synthesis Report*. Retrieved from <https://www.ipcc.ch/report/ar5/syr/>
- IPCC. (2018). *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty*.
- Lindner, M., Nikinmaa, L., Brang, P., Cantarello, E., Seidl, R. (2020). Enhancing resilience to address challenges in forest management. In F. Krumm, A. Schuck, & A. Rigling (Eds.), *How to balance forestry and biodiversity conservation. A view across Europe* (pp. 147-155). European Forest Institute; Swiss Federal Institute for Forest, Snow and Landscape Research.
- Lindner, M., Schwarz, M., Spathelf, P., de Koning, J.H.C., Jandl, R., Viszlai, I., Vanco, M. (2020). Adaptation to climate change in sustainable forest management in Europe. *FOREST EUROPE, Liaison Unit Bratislava, Zvolen*.
- Keeton, W.S., Chernyavskyy, M., Gratzner, G., Main-Korn, M., & Bihun, Y. (2010). Structural characteristics and aboveground biomass of old-growth spruce-fir stands in the eastern Carpathian Mountains, Ukraine. *Plant Biosystems* 11:1-12.
- Keeton, W.S., Franklin, J.F., & Mote, P.W. (2007). Climate variability, climate change, and western wildfire with implications for the suburban-wildland interface. Pages 223-255 in: A. Troy and R. Kennedy (eds.). *Living on the Edge: Economic, Institutional and Management Perspectives on Wildfire Hazard in the Urban Interface*. *Advances in the Economics of Environmental Resources*, Vol 6. Elsevier Sciences, New York, NY.
- Keeton, W.S., Angelstam, P., Baumflek, M., Bihun, Y., Chernyavskyy, M., Crow, S.M., Deyneka, A., Elbakidze, M., Farley, J., Kovalyshyn, V., Mahura, B., Myklush, S., Nunery, J.R., Solovity, I. & Zahvoyska, L. (2013). Sustainable forest management alternatives for the Carpathian Mountain region, with a focus on Ukraine. Pages 331-352 in J. Kozak, K. Ostapowicz, A. Bytnerowicz, and B. Wyzga (eds.) *The Carpathians: Integrating Nature and Society Towards Sustainability*. Springer-Verlag, Berlin and Heidelberg, Germany.
- Kholiavchuk, D., Gurgiser, W., & Mayr, S. 2023. Climate change in the Carpathian forests: recent developments. Presentation at the 7th Forum Carpaticum, Krakow, Poland. September 2023.
- Kobiv, Y. (2018). Trends in Population Size of Rare Plant Species in the Alpine Habitats of the Ukrainian Carpathians under Climate Change. *Diversity* 10:62 <https://doi.org/10.3390/d10030062>
- Korol, M., Havryliuk, S., Tokar, O., Gusti, M. (2022). Spatial structure, biodiversity indicators and carbon stocks of the old-growth natural forests in the protected areas of the Ukrainian Carpathians. *Environmental Sciences Proceedings* 13(1):27. <https://doi.org/10.3390/IECF2021-10803>
- Kuemmerle, T., Olofsson, P., Chaskovskyy, O., Baumann, M., Ostapowicz, K., Woodcock, C.E., Houghton, R., Hostert, P., Keeton, W.S. & Radeloff, V.C. (2011). Post-Soviet farmland abandonment, forest recovery, and carbon sequestration in western Ukraine. *Global Change Biology* 17:1335–1349.
- Kruhlov, I., Thom, D., Chaskovskyy, O., Keeton, W.S. & Scheller, R.M. (2018). Future forest landscapes of the Carpathians: vegetation and carbon dynamics under climate change. *Regional Environmental Change* 18:1555-1567.
- Mátyás, C., Berki, I., Csilléry, K., Fusco, G., Höhn, M., & Hufnagel, L. (2019). Climate change impacts and adaptation strategies in forestry: A case study in the Carpathian Mountains. *Sustainability* 11(10): 1-20.
- Mikoláš, M., Piovesan, G., Ahlström, A., Donato, D.C., Gloor, R., Hofmeister, J., Keeton, W.S., Muys, B., Sabatini, F.M., Svoboda, M. & Kuemmerle, T. (2023). Protect old-growth forests in Europe now. *Science* 380:466.
- Mikoláš, M., Svitok, M., Bače, R., Meigs, G.W., Keeton, W.S., et al. (2021). Natural disturbance impacts on trade-offs and co-benefits of forest biodiversity and carbon. *Proceedings of the Royal Society. B* 288: 20211631. <https://doi.org/10.1098/rspb.2021.1631>
- Mikoláš, M., Ujházy, K., Jasík, M., Wiezik, M., Gallay, I., Polák, P., Vysoký, J., Čiliak, M., Meigs, G. W., Svoboda, M., Trotsiuk, V., and Keeton, W. S. (2019). Primary forest distribution and representation in a Central European landscape: Results of a large-scale field-based census. *Forest Ecology and Management* 449: 117466, <https://doi.org/10.1016/j.foreco.2019.117466>.
- Millar, C.I., Stephenson, N.L. & Stephens, S.L. (2007). "Climate change and forests of the future: managing in the face of uncertainty." *Ecological Applications* 17: 2145–2151.
- Nabuurs, G.-J., Verkerk, P.J., Schelhaas, M.-J., Olabarria, J.R.G., Trasobares, A. & Cienfiala, E. (2018). Climate-Smart Forestry: mitigation impacts in three European regions. *From Science to Policy* 6. European Forest Institute.
- Panayotov, A. Ruete, B. Simovski, J. Stillhard, J. Svensson, J. Szwagrzyk, O.P. Tikkanen, K. Vandekerckhove, R. Volosyanchuk, T. Vrska, T. Zlatanov, T. Kuemmerle. (2020). Protection gaps and restoration opportunities for primary forests in Europe. *Diversity and Distributions* 00: 1-17. <https://doi.org/10.1111/ddi.13158>.

- Penuelas, J., Fernández-Martínez, M., Vallicrosa, H. et al. (2020). Increasing atmospheric CO₂ concentrations correlate with declining nutritional status of European forests. *Nature Communications: Biology* 3: 125. <https://doi.org/10.1038/s42003-020-0839-y>
- Petit-Cailleux, C., Davi, H., Lefèvre, F., Verkerk, P.J., Fady, B., Lindner, M., Oddou-Muratorio S. (2021). Tree Mortality Risks Under Climate Change in Europe: Assessment of Silviculture Practices and Genetic Conservation Networks. *Frontiers in Ecology and Evolution* 9: 706414.
- Prvălie, R., Sirodoev, I., Nita, I.A., Patriche, C., Dumitrașcu, M., Roșca, B., Tișcovschi, A., Bandoc, G., Săvulescu, I., Mănoiu, V. & Birsan, M.-V. 2022. NDVI-based ecological dynamics of forest vegetation and its relationship to climate change in Romania during 1987–2018. *Ecological Indicators* 136: 108629. <https://doi.org/10.1016/j.ecolind.2022.108629>.
- Pretzsch, H., Forrester, D.I. & Bauhus, J. (2017). *Mixed-Species Forest: Ecology and Management* 653. Germany: Springer-Verlag.
- Sabatini, F.M., Burrascano, S., Keeton, W.S., et al. (2018). Where are Europe's last primary forests? *Diversity and Distributions* 24:1426-1439. https://www.researchgate.net/publication/325360584_Where_are_Europe's_last_primary_forests.
- Sabatini, F.M., Keeton, W.S., Lindner, M., Svoboda, M., Verkerk, P.J., Bauhus, J., Bruelheide, H., Burrascano, S., Debaive, N., Duarte, I., Garbarino, M., Grigoriadis, N., Lombardi, F., Mikoláš, M., Meyer, P., Motta, R., Mozgeris, G., Nunes, L., Ódor, P., Panayotov, M., Ruete, A., Simovski, B., Stillhard, J., Svensson, J., Szwagrzyk, J., Tikkanen, O.P., Vandekerhove, K., Volosyanchuk, R., Vrska, T., Zlatanov, T., Kuemmerle, T. (2020). Protection gaps and restoration opportunities for primary forests in Europe. *Diversity and Distributions* 00: 1-17 <https://doi.org/10.1111/ddi.13158>.
- Sabatini, F. M., Bluhm, H., Kun, Z., Aksenov, D., Atauri, J. A., Buchwald, E., Burrascano, S., Cateau, E., Diku, A., Duarte, I. M., López, Á. B. F., Garbarino, M., Grigoriadis, N., Horváth, F., Keren, S., Kitenberga, M., Kiš, A., Kraut, A., Ibisch, P. L., Larrieu, L., Lombardi, F., Matovic, B., Melu, R. N., Meyer, P., Midteng, R., Mikac, S., Mikoláš, M., Mozgeris, G., Panayotov, M., Pisek, R., Nunes, L., Ruete, A., Schickhofer, M., Simovski, B., Stillhard, J., Stojanovic, D., Szwagrzyk, J., Tikkanen, O.-P., Toromani, E., Volosyanchuk, R., Vrška, T., Waldherr, M., Yermokhin, M., Zlatanov, T., Zagidullina, A., & Kuemmerle, T. (2021). European primary forest database v2.0. *Nature: Scientific Data* 8: 220. <https://doi.org/10.1038/s41597-021-00988-7>.
- Science for Environment Policy. (2021). European forests for biodiversity, climate change mitigation and adaptation. *Future Brief* 25. Brief produced for the European Commission.
- Seidl, R., Rammer, W., Scheller, R. M. & Spies, T. A. (2012). An individual-based process model to simulate landscape-scale forest ecosystem dynamics, *Ecological Modelling* 231: 87–100.
- Seidl, R., M.-J. Schelhaas, W. Rammer, and P. J. Verkerk. (2014). Increasing forest disturbances in Europe and their impact on carbon storage. *Nature Climate Change* 4:806–810.
- Seidl, R., Thom, D., Kautz, M. et al. (2017). Forest disturbances under climate change. *Nature Climate Change* 7: 395–402. <https://doi.org/10.1038/nclimate3303>
- Senf, C. & Seidl, R. (2021a). Mapping the forest disturbance regimes of Europe. *Nature Sustainability* 4: 63–70. <https://doi.org/10.1038/s41893-020-00609-y>.
- Senf, C. & Seidl, R. (2021b). Persistent impacts of the 2018 drought on forest disturbance regimes in Europe. *Biogeosciences*, 18: 5223–5230. <https://doi.org/10.5194/bg-2021->
- Spinoni, J. and the CARPATCLIM project team (39 authors). (2014). Climate of the Carpathian Region in 1961-2010: Climatologies and Trends of Ten Variables. *International Journal Climatology*. <https://doi.org/10.1002/joc.4059>.
- Stoyko, S.M. (1998). Virgin ecosystems of the Carpathians and their significance for biological diversity conservation and maintenance of the sustainable development of forestry. In: *Issues of sustainable development in the Carpathian region*, vol 2, Rakhiv.
- Thom, D., Ammer, C., Annighöfer, Aszalós, P.R., Dittrich, S., Hagge, J., Keeton, W.S., Kovacs, B., Krautkrämer, O., Müller, J., von Oheimb, G. & Seidl, R. (2023). Regeneration in European beech forests after drought: the effects of microclimate, deadwood and browsing. *European Journal of Forest Research* 142: 259–273.
- Thom, D., Rammer, W. & Seidl, R. (2017). Disturbances catalyze the adaptation of forest ecosystems to changing climate conditions. *Global Change Biology*. 23: 269–282.
- Thom, D. & Seidl, R. (2016). Natural Disturbance Impacts on Ecosystem Services and Biodiversity in Temperate and Boreal Forests. *Biological Reviews* 91: 760–81.
- Thuiller, W., Lavorel S., Araújo, M.B., Sykes, M.T., Prentice, I.C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America* 102:8245-8250.
- Vitasse, Y., Bottero, A., Rebetez, M., Conedera, M., & Augustin, S. (2019). Understanding tree responses to global warming through tree-ring analysis. *Annals of Forest Science* 76: 1-11.
- Werners, S.E., Bos, E., Civic, K., Hlásny, T., Hulea, O., Jones-Walters, L., Kópataki, É., Kovbasko, A., Moors, E., Nieuwenhuis, D., van de Velde, I., Henk Zingstra, H. & Zsuffa, I. (2014). Climate change vulnerability and ecosystem-based adaptation measures in the Carpathian region: Final Report - Integrated assessment of vulnerability of environmental resources and ecosystem-based adaptation measures. Wageningen, Alterra Wageningen UR (University & Research Centre), Alterra report 2572. 132 pp.; 37 fig.; 14 tab.; 111 ref.
- Werners, S.E., Szalai, S., Kópataki, É., Kondor, A.C., Musco, E., Koch, H., Zsuffa, I., Ji. Trombik, Kuras, K., Koeck, M., Lakatos, M., Peters, R., Lambert, S. & Hlásny, T. (2014). Future imperfect: climate change and adaptation in the Carpathians. CARPATCLIM, CARPIVIA and Carpathian Convention, Vienna, Austria. 37 pp.
- Werners, S.E., Szalai, S., Zingstra, H., Kópataki, É., Beckmann, A., Bos, E., ... Zsuffa, I. (2016). Climate change adaptation in the Carpathian Mountain Region. In N. Salzmann, C. Huggel, S. U. Nussbaumer, & G. Ziervogel (Eds.), *Climate change adaptation strategies—An upstream-downstream perspective* (pp. 79–99). Heidelberg, Germany: Springer International Publishing. https://doi.org/10.1007/978-3-319-40773-9_5.
- Zamora-Camacho, F.J., Castro, J. & Gomez-Aparicio, L. (2019). Forest management and global change: A review from the Spanish perspective. *Forests* 10(5): 1-16.
- Zang, C., Hartl-Meier, C., Dittmar, C., Esper, J., Göttlein, A. & Rothe, A. (2019). Wide-spread decline of Norway spruce forests in the German mountain ranges since the late 1970s. *Forest Ecology and Management* 432: 10-18.

PHOTO CREDITS

i	Tomáš Hulík	35	Hungarian National Tourism Agency MUT
ii	Robert Balog	36	Tomáš Hulík
iv	Mircea Verghelet	37	Arek Socha
v	Tomáš Hulík	39	Ewelina Zajac (top photo)
vi	Klaudia Kuraś	39	Jaroslav Košťál (bottom photo)
vii	Mircea Verghelet	40	Zbigniew Niewiadomski
1	Wikimedia Commons	41	William Keeton
3	Zbigniew Niewiadomski	42	Juraj Vysokí
4	Montenegrin Eden	43	Ivan Kluüiar
5	Mircea Verghelet	44	Ewelina Zajac (top photo)
6	Wojciech Hurkala	44	Mircea Verghelet (bottom photo)
12	Mircea Verghelet	46	Ewelina Zajac
13	Mircea Verghelet	47	Ulrich Rueger
15	William Keeton	48-49	Mircea Verghelet
16	Miloslav Brandt (top photo)	50	Klaudia Kuraś
16	Juraj Vysokí (bottom photo)	51	Mircea Verghelet
17	Mircea Verghelet	66	Stefan Rankovic
18	Arek Socha		
19	Jaroslav Košťál		
20	Mircea Verghelet		
22	Klaudia Kuraś (top photo)		
22	Nadine (bottom photo)		
24	Arek Socha		
25	Arek Socha		
28	Ed Pylypenko		
30	Arek Socha		
31	Wikimedia Commons		
32	Mircea Verghelet		
33	Ewelina Zajac		
34	Miloslav Brandt		

ANNEX 1:

NOMINATED EXPERTS SUPPORTING THE ASSESSMENT

Expert Group for the development of the assessment of climate risks and adaptation options for Carpathian forest ecosystems and their services

The below experts were nominated by the Carpathian Convention Parties based on the CC NOTIFICATION 2021 - 7 – Requesting nomination of experts to be involved in the climate change assessment / special session at the Forum Carpaticum 2022 /Working Group Climate Change + Working Group Forest.

CZECH REPUBLIC

Miroslav Svoboda

Czech University of Life Sciences Prague

Eliška Rolfova

Ministry of the Environment of the Czech Republic

Radek Pokorný

Mendel University in Brno

HUNGARY

Borbála Galos

University of Sopron

Imelda Somodi

Centre for Ecological Research

POLAND

Bożydar Neroj

Bureau for Forest Management and Geodesy

Wojciech Grodzki

Forest Research Institute

Małgorzata Czyżewska

Directorate General of the State Forest of Poland

ROMANIA

Laurentiu Radu

Ministry of Environment, Waters and Forest

Liliana Virtopeanu

Ministry of Environment, Waters and Forest of Romania

Borz Stelian Alexandru

Transilvania University of Brasov, Department of Forest Engineering

Păcurar Victor Dan

Transilvania University of Brasov

Sorin Cheval

National Meteorological Administration of Romania

Popa Ionel

Forest Research and Management Institute, Romania

SLOVAKIA

Libor Ulrych

State Nature Conservancy of Slovak Republic

SERBIA

Ilija Dordevic

Institute of forestry, Department for spatial planning, GIS and forest policy, Assistant director for international cooperation

UKRAINE

Liubov Poliakova

Head of International Cooperation, Science and Public Relation Division, State Forest Resources Agency

Volodymyr Korzhov

Deputy Head of Ukrainian Scientific Institute of Mountain Forestry

COORDINATORS

William Keeton

University of Vermont and Member of the Science for the Carpathians

Sabine McCallum

Associate Programme Management Officer, UN Environment Programme-Ecosystems Division

Klaudia Kuraś

Carpathian Convention Coordination Expert, UN Environment Programme-Secretariat of the Carpathian Convention

ANNEX 2: SURVEY

INTRODUCTION

The Carpathian Convention Conference of the Parties at its 6th meeting (COP6, 2020) through its decisions³² encouraged the development of an assessment of the impacts of climate change on the Carpathian forests and their ecosystems services by relevant Convention Working Groups and partners and with support of the Convention Secretariat. Subsequently, this activity has been included in the Implementation Framework 2030 accompanying the Long-term Vision towards combating climate change in the Carpathians.

This survey aims to gather information for developing the assessment of climate change risks and impacts on Carpathian Forest ecosystems and their services along a draft table of contents (see Annex) that has been presented and agreed at the 1st Expert Workshop on 16 November 2021 (online).

There are 4 sections of the survey:

A: CONTACT and CONTRIBUTORS

B: KNOWLEDGE BASE on CLIMATE CHANGE RISKS and IMPACTS on CARPATHIAN FOREST ECOSYSTEMS and their services

C: PRACTICAL EXAMPLES / CASE STUDIES

D: REFERENCES

The Secretariat of the Carpathian Convention together with Dr. William Keeton, University of Vermont and Member of the Science for the Carpathians, highly appreciates your willingness to contribute to assessment by sharing your insights and expertise through this survey. We would also encourage you to consult with national colleagues for further contributions.

Please return the filled in survey until 28.01.2022. Many thanks in advance for your valuable inputs!

SECTION A. CONTACT AND CONTRIBUTORS

* 1. Contact details

Name:

Click or tap here to enter text.

Institution you represent:

Click or tap here to enter text.

Type of institution:

Choose an item.

If you chose Other, please specify below:

Click or tap here to enter text.

Gender:

Choose an item.

Email Address:

Click or tap here to enter text.

* 2. Country

This survey will ask you a series of questions about a particular country where you operate. We appreciate that you may work in multiple locations, so please indicate one below that you will discuss here.

Please use the dropdown list below to select the country:

Choose an item.

* 3. CONTRIBUTORS

You may want to list colleagues that contributed to filling in this survey and shall be acknowledged:

Name:

Click or tap here to enter text.

Institution

Click or tap here to enter text.

SECTION B. KNOWLEDGE BASE ON CLIMATE CHANGE RISKS AND IMPACTS ON CARPATHIAN FOREST ECOSYSTEMS AND THEIR SERVICES

* 1. Key risks and impacts

For distinguishing between risk and impacts, we are using the concepts of [how the IPCC assesses and communicates to decision-makers](#):

The 'core' definition of **risk** is “**the potential for adverse consequences**”:

- The word “potential” makes clear that uncertainty, or more broadly, incomplete knowledge (as defined in IPCC), is a key element of the concept of risk.
- In IPCC use, risk refers only to negative (“adverse”) consequences³³.

The term **impact** is used to describe the consequences of realised risks on natural and human systems, where risks result from the interactions of climate-related hazards (including extreme weather and climate events), exposure, and vulnerability.

- Impacts may be referred to as consequences or outcomes occurring within a specific time period.
- Impacts can be adverse or beneficial.

From your experience, please indicate the main risks and potential impacts along the identified key topics:

FOREST GROWTH AND PRODUCTIVITY

Key risk	Potential impacts (consequences, outcomes)

BIOMASS AND CARBON STOCKS

Key risk	Potential impacts (consequences, outcomes)

TREE MORTALITY

Key risk	Potential impacts (consequences, outcomes)

CHANGES IN SPECIES RANGE, HABITAT SHIFTS AND ABUNDANCE

Key risk	Potential impacts (consequences, outcomes)

INVASION BY NON-NATIVE SPECIES

Key risk	Potential impacts (consequences, outcomes)

FOREST ECOSYSTEM SERVICES

Key risk	Potential impacts (consequences, outcomes)

FOREST – WATER INTERACTIONS, INCLUDING HYDROLOGIC REGULATION AND RIPARIAN DYNAMICS

Key risk	Potential impacts (consequences, outcomes)

*** 2. Adaptation responses**

From the key risks and impacts identified under B.1, please indicate adaptation response options that you are aware of and briefly highlight their intended effects for each key topic. Please note that there is an additional possibility to share adaptation response options that are cross-cutting in tackling more than one of the key topics.

FOREST GROWTH AND PRODUCTIVITY

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

BIOMASS AND CARBON STOCKS

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

TREE MORTALITY

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

CHANGES IN SPECIES RANGE, HABITAT SHIFTS AND ABUNDANCE

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

INVASION BY NON-NATIVE SPECIES

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

FOREST ECOSYSTEM SERVICES

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

FOREST – WATER INTERACTIONS, INCLUDING HYDROLOGIC REGULATION AND RIPARIAN DYNAMICS

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

CROSS-CUTTING

Please indicate adaptation response options addressing the key risks and impacts mentioned above:

Name / Key word	Main Impact/Risk addressed	Brief description	Intended effect	Pros and cons (if any)

ADDITIONAL FOCUSED QUESTIONS DERIVED FROM THE EXPERT DISCUSSION

Please provide your thoughts on the following topics regarding specific adaptation response options raised at our first Expert Workshop on 16 November 2021. In case you already covered one or more of these additional questions, please refer to the respective section above.

Planting and management of exotic species.

Should use of exotic, non-European species comprise an element of adaptive management? Where, when, and how?

Click or tap here to enter text.

Role of landscape level planning, including a diversity of forest zonation and management strategies.

What is your view on the role of protected areas vs. active adaptive management?

Click or tap here to enter text.

Expanded use of “close-to-nature” silviculture (e.g., selection harvesting, continuous cover forestry, retention forestry, etc.).

How is the forest sector in your country considering broadening its portfolio of forest management practices to adapt to climate change, including altered disturbance regimes?

Click or tap here to enter text.

Forest road density, design, and location.

How should forest road systems be managed to reduce vulnerabilities to flooding?

Click or tap here to enter text.

Long-term adaptive forest management objectives.

Should we manage for the historic, current, or future potential vegetation? How is the forest sector in your country approaching these challenging questions?

Click or tap here to enter text.

Public policy, perception, and science.

What are the greatest challenges you face relating to formulating adaptation responses, given the interplay between public perception and public policy that may or may not always be consistent with the science?

Click or tap here to enter text.

Forest harvest rotations.

Is the forest sector in your country considering reducing or increasing forest harvest rotations? Why or why not?

Click or tap here to enter text.

Adaptation to altered natural disturbance regimes.

How is the forest sector in your country adapting to increasing risks of bark beetles, wind, fire, and drought?

Click or tap here to enter text.

Mix of old vs. younger forest stands.

How is the forest sector in your country adjusting the mix of forest ages as adaptation to disturbance risk, for the purpose of carbon management, or to conserve biodiversity in the face of climate change?

Click or tap here to enter text.

* 3. Further opportunities and pathways

If you are aware of any further unused potentials and opportunities for effective adaptation responses / pathways (e.g., in other countries/mountain regions), please briefly describe:

Click or tap here to enter text.

Please specifically outline possible approaches and pathways you know focusing on inclusive ecosystem restoration using Nature based Solutions (NbS) and Ecosystem based Adaptation (EbA):

Click or tap here to enter text.

* 4. Key initiatives

Please share ongoing relevant initiatives / larger scale projects in your country / the Carpathian region / elsewhere:

IN YOUR COUNTRY

Name of the initiative /project	Duration of implementation	Brief description	Weblink (if available)	Contact for further information

IN THE CARPATHIAN REGION

Name of the initiative /project	Duration of implementation	Brief description	Weblink (if available)	Contact for further information

ELSEWHERE

Name of the initiative /project	Duration of implementation	Brief description	Weblink (if available)	Contact for further information

* 5. Limitation and barriers to overcome

From your experience, please highlight the most common limitations and barriers to overcome for developing and implementing effective adaptation responses. These could potentially link to policy frameworks, shortcomings with financing for implementation, cross-border cooperation, etc.

Click or tap here to enter text.

* 6. Knowledge gaps and research needs

In your opinion, where do we still have major knowledge gaps and thus research needs toward better informed decision-making for forest ecosystem climate change adaptation?

Please indicate in which areas you see knowledge gaps and research needs and briefly explain why:

Knowledge gap	Research need	Brief explanation

SECTION C. PRACTICAL EXAMPLES / CASE STUDIES

Within this section we aim to collect practical examples that could be showcased as promising approaches with upscaling potential to other countries/regions.

As a general orientation for considering practical examples to share, please reflect the following aspects¹:

KEY DIMENSIONS TO QUALITY FOR A MOUNTAIN ADAPTATION SOLUTION

Relevant

The solution addresses one or more current or anticipated mountain-specific climate change risks and provides a promising approach to becoming effective in tackling the issue at stake. In this regard, the solution is based on scientific evidence and/or traditional knowledge and practices.

Practical and feasible

The solution can be implemented on relevant timescales to address the risks in question, is realistic in terms of resources available (human and financial) and tailored to the actors and their capacities needed for implementation and is sustainable in the longer term (both human capacities and financial resources can realistically be maintained).

Direct benefits and co-benefits

The solution promotes ecological, economic and/or social benefits. It shows synergy with and offers co-benefits to climate change mitigation and other sustainable mountain development topics, such as eradication of poverty, averting unemployment, provision of humanitarian aid in case of conflict or disasters, universal health coverage and education, achieving gender equality and empowering women and girls.

Flexible and robust

The solution is designed in a way that allows for adjustments and incremental implementation and reiteration depending on the level and degree of climate change, i.e. allows for adaptive management and responds to multiple interests and purposes. Thus, the solution is robust in terms of maintaining its effectiveness under a range of different climatic and socio-economic development scenarios. In doing so, the solution should ideally have built-in mechanisms to enable its monitoring and evaluation of time.

Replicable and/or scalable

The solution including its enabling factors has the potential for adjustment, replication or upscaling in other geographic, social or sectorial contexts (even though as such customized and tailored to specific local circumstances).

Legitimate and coherent

The solution is politically, culturally, and socially accepted. The solution is not in conflict with other adaptation or sustainable development efforts and coherent with existing or planned policies on local, regional and national level (please note that this does not translate into a requirement that the solution is already integrated in a local, regional or national policy!)

1. These key dimensions are being used under the Adaptation at Altitude programme for gathering and selecting mountain adaptation solutions in South Caucasus and East Africa.

Please share one or more promising adaptation approaches from your country that are already in the process of being implemented describing them along the following simple structure. For sharing more than one case study please simply copy the text box below:

Name of the practical example/case study:

Click or tap here to enter text.

Description:

- **The issue**
Short description of the issue to be tackled, which specific related risk/s and impacts are or were being addressed and what the evidence base/need for developing this adaptation response in this particular area is [responding to the dimension of Relevance].
Click or tap here to enter text.
- **The solution**
Detailed description of the solution in response to the issue at stake: Short background why the approach has been chosen for addressing the specific mountain-related risk/s and how it has been designed for effective implementation in the geographical location. If applicable, reference shall be made to necessary enabling factors that contribute to the solution's success such as social inclusion, women empowerment, taking account of the broader socio-economic context, securing political commitment and financing. The description shall also include the time planned for implementing the solution in this area, built-in mechanisms for evaluation and feedback loops, room for adjustment if needed and the timescale for which the solution is planned to last [responding to the dimensions of Practical and efficient; Flexible and robust].
Click or tap here to enter text.
- **Coverage and Impact**
Brief summary of the main effects adverse which are already evident through implementing the solution, including reference to all areas where the approach provides impacts at the moment of writing the text. A portrait of a beneficiary or a project "owner" will give a more personal aspect to the text. The text can be supported by further elements such as graphs and photographs to better explain the impact of the solution [responding to the dimensions of Direct benefits and co-benefits; Legitimate and coherent].
Click or tap here to enter text.
- **Applicability**
Short description about the potential to upscale, replicate or transfer this solution in a different context or continent. In particular, the context specificity but also characteristics of the location are relevant factors for the transferability and scalability of a solution. These factors include, inter alia, the social and cultural context in which the solution is implemented (e.g. the solution responds to gender-differentiated vulnerabilities, is socially accepted and generally compatible with mountain livelihoods systems), characteristics of beneficiaries of the solutions (e.g. in terms of risk aversion, capacity building towards increasing climate resilience), possibly technology characteristics (e.g. costs, familiarity, perceived usefulness, profitability, co-benefits and/or 'no regrets' potential), the policy environment and other transfer mechanisms such as incentives. Referring to the specific location the solution is being implemented, similar climatic and geophysical factors may also be a precondition for a solution to achieve a similar outcome in a different geographical region [responding to the dimension of Replicable and/or scalable].
Click or tap here to enter text.

SECTION D. REFERENCES AND ADDITIONAL INFORMATION

* 1. References and further information

Please use this section to share further references and additional information that you see relevant for the assessment (from your country / from the Carpathian region / from elsewhere).

FROM YOUR COUNTRY

Click or tap here to enter text.

FROM THE CARPATHIAN REGION

Click or tap here to enter text.

FROM ELSEWHERE

Click or tap here to enter text.

* 2. Final comments

Please add any other comments or thoughts you would like to share here regarding climate change vulnerabilities and adaptation in forest ecosystems in your country or for your institution.

Click or tap here to enter text.

